



Short Communication

Advancements in Life Sciences – International Quarterly Journal of Biological Sciences

ARTICLE INFO

Open Access



Date Received:
30/05/2022;
Date Revised:
29/01/2023;
Date Published Online:
30/06/2023;

Molecular phylogeny and morphological characterization of *Leucocoprinus birnbaumii* from Punjab, Pakistan

Hira Ijaz¹, Maham Mureed¹, Aneela Yasmeen², Arooj Naseer³, Abrar Hussain¹, Sana Jabeen^{1*}

Authors' Affiliation:

1. Department of Botany, Division of Science and Technology, University of Education, Township, Lahore - Pakistan
2. Centre of Excellence in Molecular Biology, University of the Punjab, Lahore - Pakistan
3. Institute of Botany, University of the Punjab, Quaid-e-Azam Campus, Lahore - Pakistan

*Corresponding Author:

Sana Jabeen
Email:
sanajabeen@ue.edu.pk

How to Cite:

Ijaz H, Mureed M, Yasmeen A, Naseer A, Hussain A, Jabeen S (2023). Molecular phylogeny and morphological characterization of *Leucocoprinus birnbaumii* from Punjab, Pakistan. Adv. Life Sci. 10(2): 148-151.

Keywords:

Humid subtropical climate; ITS; nrDNA; Taxonomy

Abstract

Background: *Leucocoprinus* in *Agaricaceae* is a genus distributed worldwide and represented by nearly 50 species. From Pakistan, only four species in this genus have been known so far.

Methods: The specimen was identified by morphological and anatomical characters combining with sequence analysis of nrDNA-ITS region.

Results: *Leucocoprinus birnbaumii*, a bright yellow mushroom found on wood log from Sialkot district, Punjab, Pakistan. This species is identified by its medium sized bell-shaped pileus bearing the same colored squamule with sulcate striate margins, presence of metachromatic basidiospores and pseudoparaphyses around the basidia. Molecular data also supported its taxonomy as *L. birnbaumii*. The species is being described for the first time based on morphological and anatomical characters in combination with molecular phylogeny using ITS region of nrDNA.

Conclusion: *Leucocoprinus birnbaumii* is widely distributed in Asia, Africa, Europe, North America and South America. It grows in tropical and sub-tropical areas. Its occurrence in Sialkot, Punjab, Pakistan with humid subtropical climate represents its wide ecological amplitude and geographic distribution.



Introduction

Leucocoprinus Pat. is a genus of lepiotaceous fungi in *Agaricaceae* distributed worldwide in the tropical and sub-tropical areas [1]. This genus is represented by nearly 50 species and typified by *L. cepistipes* (Sowerby) Pat [2-4]. It occurs naturally in Asia, Central and Southern Africa, Europe, North America and South America and mostly common in hot, humid weather [4-9]. From Pakistan, only four species in this genus have been known so far in the form of a list with material examined but their detailed descriptions are not given [10]. The members of the genus are saprobic and quickly grow in potting soils, greenhouses and man-made other organic rich matters [6, 9, 11, 12].

Leucocoprinus is characterized by pale, white, metachromatic basidiospores and the sulcate, often plicate pileus margins. These characters are intermediate between *Leucoagaricus* Locq. ex Singer and *Macrolepiota* Singer [13]. This difference was clarified by the discovery of presence of pseudoparaphyses between its basidia that are absent in *Leucoagaricus* species. Clamp connections are absent in all the tissues of *Leucocoprinus* [9, 11, 14].

The current study emphasizes on characterization of specimens found morphologically similar to *Leucocoprinus* growing on wooden logs in Sialkot district, Punjab, Pakistan. The detailed morphological and anatomical description and molecular phylogenetic analysis based on ITS region of nrDNA was carried out to identify the specimens up to species level.

Methods

The specimens were collected from Khajoriwala, Sialkot district, Gujranwala division, Punjab, Pakistan during the monsoon season of July 2019 and their macroscopic characters were noted. Colors were noted and codes were given according to the Munsell (1975) color chart [15]. Photographs were taken and then basidiomata were subjected to air dry for preservation. Microscopic structures were studied under the microscope (Xsz 107BN) with a 100× objective lens. All microscopic features including size, shape, color, and arrangements of hyphae in different tissues were noted. Measurements were recorded using Motic Images Plus 2.0. For the measurements of basidiospores, the notation [s/b/c] was used, where s represents the number of basidiospores, measured from b basidiomata and c collections. Extreme values of basidiospores dimensions (l × w) were given in parenthesis. Q values were calculated as l × w ratio. Q values for spores were interpreted following Bas (1969) [16]. Drawings were drawn from the laptop screen. The examined specimens were deposited in the herbarium of University of Education, Lahore, Pakistan (UEH).

For genomic DNA Extract-N-Amp™ kit was used following the manufacturer's instructions. The ITS region was amplified and sequenced from the sequence service using ITS1F (forward primer) and ITS4 (reverse primer) primers. Sequences obtained were analyzed in BioEdit sequence alignment editor version 7.2.5 [17]. The consensus sequence was BLAST searched at NCBI database (nucleotide BLAST). Closely relates sequences were selected from GenBank to reconstruct phylogeny. Published sequences of the closest relatives of the species were also included in the dataset. All the sequences were aligned online using MUSCLE [18] alignment tool. The final aligned data set was analyzed through MEGA 6.0 software for reassessing the phylogeny (Figure 1). Kimura 2-parameter [19] was selected as an evolution model for maximum likelihood analysis using the best-fit substitution model approach in MEGA [20].

Results

Molecular phylogeny

The total fragment size of the target region of nrDNA of collected specimen yielded 784 base pairs. The BLAST results showed 98.56–99.46% similarity with *Leucocoprinus birnbaumii* sequences (LN827701, U85323 and OL142279) from Egypt, UK and USA. These sequences along with other sequences from GenBank were retrieved for the phylogenetic analysis. *Leucoagaricus rubrotinctus* (Peck) Singer (KP300876) and *L. lahorensiformis* Hussain *et al.*, (KU647729) were used to root the tree following Justo *et al.* (2021) phylogeny [4].

The final dataset of ITS region contains 58 nucleotide sequences including outgroups representing 731 positions. Among these positions, 438 were conserved, 285 were variable, 238 were parsimony informative and 46 were found as singletons. In phylogenetic tree, sequence.

Taxonomy

***Leucocoprinus birnbaumii* (Corda) Singer, Sydowia 15(1-6): 67 (1962)**

Description

Pileus up to 4.7 cm broad, campanulate, broadly umbonate, bright yellow (5Y8/10); fibrillose, floccose to loosely scattered concolorous squamules; margins incurved, sometimes straight, sulcate-striate, pale yellow (2.5Y8/8); context yellow (7.5Y 9/6). Lamellae free, sub-distant, ventricose, pale yellow (2.5Y8/8); edges fimbriate. Lamellulae absent. Stipe 6.5–9 × 0.4–0.9 cm; central, equal, base bulbous upto 1.4 cm, pale yellow (7.5Y 9/3); surface floccose squamulose; light yellow (5Y8/10); context yellow (7.5Y9/6). Annulus central, moveable, uplifted, thin, band like; yellow

(2.5Y8/12), leaving remnants on pileus margins. Volva absent.




Figure 1: Phylogenetic sequence analysis ITS sequences representing *Leucocoprinus birnbaumii* and related species in the genus along with two outgroup taxa. The sequence generated during this study is marked with the bullet.



Figure 2: Basidiomata of *Leucocoprinus birnbaumii* UEH-F0001 showing its morphological features. Scale bars = 1 cm.

Basidiospores [60/3/1] (10)10.1–11.7(12) × (6.9)7.2–8.3(8.4) μm, Q (1.4)1.41–1.43(1.44), avQ = 1.42, ellipsoid, amygdaliform, thick walled, apiculate with a prominent germ pore, smooth, metachromatic. Basidia (18.3)18.5–23.7(24.6) × (9.1)9.3–14(14.5) μm, cylindrical to clavate, 4-spored, thin walled, hyaline. Pseudoparaphyses (14.7)16–17(17.5) × (12.2)12.5–

14.5(15) μm, broadly clavate, thin walled, hyaline. Cheilocystidia (45)48–50.6(54) × (9.4)11.0–11.7(14) μm, clavate to narrowly utriform or fusiform, flexuous, thin walled. Pleurocystidia absent. Stipitipellis hyphae (5)5.1–7(9) μm, cylindrical, septate, thin walled. Pileipellis hyphae (5.7)8–9.3(9.6) μm, cylindrical, thin walled. Clamp connections absent.

Material Examined: Pakistan, Punjab, Gujranwala division, Sialkot district, Khajoriwala, 256 m a.s.l. from wooden roof, 25 July 2019, Maham Mureed MM34-SJ220 (UEH-F0001; GenBank accession: ON627825).




Figure 3: *Leucocoprinus birnbaumii* UEH-F0001. A. Basidiospores; B. Cheilocystidia; C. Basidia; D. Pseudoparaphyses; E. Stipitipellis; F. Pileipellis. Scale bars = 10 μm.

Discussion

Leucocoprinus birnbaumii is the best-known member of *Leucocoprinus*, previously placed in different genera; *Agaricus* L., *Bolbitius* Fr. and *Lepiota* (Pers.) Gray and was known as *Agaricus aureus* F.M. Bailey and *A. birnbaumii* Corda. which are now reduced to synonyms (IndexFungorum) [8, 21]. It is commonly known as flowerpot parasol and the plantpot dapperling as it is usually found growing in flowerpots and greenhouses [22].

Leucocoprinus birnbaumii is identified by bright yellow, medium sized basidiomata having bell shaped pileus bearing same colored squamules with sulcate striate margins, presence of metachromatic basidiospores and pseudoparaphyses around the basidia [5-9, 13]. It is an inedible, poisonous mushroom widely distributed in Asia, Africa, Europe and North and South America and grow on decayed plant matter, humus rich or composted soil in flowerpots and flowerboxes in green houses or in buildings in tropical and sub-tropical areas [1, 6, 8, 9, 11]. A series of alkaloids and fatty acids have been studied and

identified in *L. birnbaumii* which play an important role against the bacterial activity [9, 22, 23].

The present collection was found solitary on wooden log from the humid subtropical climate (Köppen climate classification) of Gujranwala division, Sialkot district, Punjab, Pakistan. The morphological and anatomical descriptions and molecular analysis of the *L. birnbaumii* provides a baseline data for future taxonomic work on this species and related taxa which will be helpful in various applied fields particularly pharmaceutical industry.

Competing Interest

The authors declare that there is no conflict of interest.

Author Contributions

Hira Ijaz: Drafted the manuscript and helped in phylogenetic analysis

Maham Mureed: Collected the sample and worked on it for her thesis

Aneela Yasmeen: Helped in molecular work and formatted the document

Arooj Naseer: Helped in microscopy

Abrar Hussain: Supervisor of Maham Mureed, reviewed the document.

Sana Jabeen: Co supervisor of Maham Mureed, took responsibility of the document, reviewed it, made plates and did phylogenetic analysis.

References

- Vellinga EC. Ecology and distribution of lepiotaceous fungi-a review. *Nova Hedwigia*, (2004); 78: 273–299.
- Kirk PM, Cannon PF, Minter DW, Stalpers JA. *Dictionary of the Fungi* (10th ed.). (2008); p.374. Wallingford, UK: CAB International,
- de Albuquerque MP, de Carvalho VF. *Leucocoprinus fluminensis* (*Agaricaceae, Basidiomycota*), a new species from southwest Brazilian Rain Forest. *Neotropical Biology and Conservation*, (2012); 7(3): 158–161.
- Justo A, Angelini C, Bizzi A. The genera *Leucoagaricus* and *Leucocoprinus* in the Dominican Republic. *Mycologia*, (2021); 113(2): 348–389.
- Reid DA, Eicker A. South African fungi. 2. Some species of *Leucoagaricus* and *Leucocoprinus*. *South African Journal of Botany*, (1993); 59(1), 85–97.
- Birkebak JM. The genus *Leucocoprinus* in western Washington. *Mycotaxon*, (2010); 112(1): 83–102.

- Wojewoda W, Karasiński D. Invasive macrofungi (*Ascomycota* and *Basidiomycota*) in Poland. *Biological invasions in Poland*, (2010); 1(7).
- Verma RK, Pandro V. Diversity of macro-fungi in Central India-XIII: *Leucocoprinus badhamii* and *Leucocoprinus birnbaumii*. *Van Sangyan*, (2018); 5(5&6): 28–37.
- El-Fallal A, El-Sayed AK, El-Gharabawy HM. *Podaxis pistillar* (L.) Fr. and *Leucocoprinus birnbaumii* (Corda) Singer; new addition to macrofungi of Egypt. *Egyptian Journal of Botany*, (2019); 59(2): 413–423.
- Ahmad S, Iqbal SH, Khalid AN. *Fungi of Pakistan*. (1997); Sultan Ahmad Mycological Society of Pakistan.
- Vellinga EC. *Leucocoprinus*. In Noorderloos M. E., Kuyper Th. W., Vellinga E. C. (eds). *Flora Agaricina Neerlandica*; (2001); AA Balkema, Rotterdam, The Netherlands.
- Kumaresan V, Veeramohan R, Ganesan T. Diversity of white-spored agarics in Puducherry. *Journal of Mycology and Plant Pathology*. (2011); 41: 518–523.
- Singer R. *The Agaricales in modern taxonomy*. 4th ed. (1986.); 982p. Koeltz Scientific Books, Koenigstein,
- Vellinga EC. Genera in the family *Agaricaceae*-Evidence from nrITS and nrLSU sequences. *Mycological Research* (2004); 108: 354–377.
- Munsell AH. *Munsell Soil Color Charts*. Macbeth Division of Kollmorgen Corporation. (1975). Baltimore, Maryland.
- Bas C. Morphology and subdivision of *Amanita* and a monograph of its section *Lepidella*. *Persoonia*, (1969); 5: 96–97.
- Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acid Series*, (1999); 41: 95–98.
- Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, (2004); 32(5): 1792–1797.
- Kimura M. A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*, (1980); 16: 111–120.
- Tamura K, Stecher G, Peterson D, Filipksi A, Kumar S. MEGA6: Molecular Evolutionary Genetics Analysis Version 6.0. *Molecular Biology and Evolution*, (2013); 30(12): 2725–2729.
- Roberts P, Evans S. *The Book of Fungi*. (2011). Chicago, USA: University of Chicago Press.
- Brkljača R, Urban S. Rapid dereplication and identification of the bioactive constituents from the fungus, *Leucocoprinus birnbaumii*. *Natural Product Communications*, (2015); 10: 95–98.
- Vellinga EC. Nomenclatural overview of Lepiotaceous fungi. Version 4.4 2/4/2009. http://plantbio.berkeley.edu/~bruns/ev/vellinga_nomencl_v47_feb2009.pdf (Accessed 5/19/2009).



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License. To read the copy of this license please visit: <https://creativecommons.org/licenses/by-nc/4.0/>