

ARTICLE INFO

Open Access



Date Received:

30/07/2023;

Date Revised:

31/08/2023;

Date Published Online:

31/12/2023;

Date Updated:

09/09/2025

Authors' Affiliation:

1. Department of Biology,
Faculty of Science, Samtah,
Jazan University - Saudi Arabia

2. Department of Zoology,
Faculty of Science, Assiut
University, Assiut - Egypt

3. Department of Botany and
Microbiology, Faculty of
Science, Assiut - Egypt

4. Pharmacognosy Department,
College of Pharmacy, King Saud
University, Riyadh - Saudi
Arabia

5. Pharmacognosy Department,
Faculty of Pharmacy, Assiut
University, Assiut - Egypt

6. Department of Science
Laboratories, College of Science
and Arts, Qassim University, Ar
Rass - Saudi Arabia

*Corresponding Authors:

Ahmed. M. H. Ali

Email:

shafa2000us@yahoo.com

How to Cite:

Mohamed NH, Ali AHM,
Tawfik AI, Ismail MA,
Mageed WMA, Shoreit AAM
(2023). Feeding deterrence
and larvicidal effects of latex
serum and latex-synthesized
nanoparticles of *Calotropis*
procera against the cotton
leafworm, *Spodoptera*
littoralis. Adv. Life Sci. 10(4):
585-592.

Keywords:

Calotropis; Latex;
Nanoparticle; Spodoptera;
Antifeedant activity

Editorial Note:

You are viewing the latest
version of this article having
language corrections.

Feeding deterrence and larvicidal effects of latex serum and latex-synthesized nanoparticles of *Calotropis procera* against the cotton leafworm, *Spodoptera littoralis*

Nadia. H. Mohamed¹, Ahmed. M. H. Ali^{2,6,*}, Amer I. Tawfik², Mady A. Ismail³, Wael. M. Abdel-Mageed^{4,5}, Ahmed A. M. Shoreit³

Abstract

Background: *Spodoptera littoralis* is an important agricultural pest; thus, knowledge about the effect of latex serum and latex-synthesized nanoparticles of *Calotropis procera* on this species can assist in its management. Nanotechnology is currently taking root in the agriculture economy as a substitute for pest management that is targeted, safe, and effective. The repellent and antifeedant efficacy of *C. procera* latex serum and its nanoparticles against *S. littoralis* fourth instar larvae was investigated in this work.

Methods: The process of synthesizing silver nanoparticles has been carried out, followed by their subsequent characterization. The objective of this study was to assess the antifeedant properties of the latex serum and its nanoparticles against the 4th instar larvae of *S. littoralis* through the implementation of a bioassay.

Results: With rising latex serum content, the proportion of repellency and starvation increased. In comparison to insects injected with latex serum, utilizing nanoparticles of LAgNPs via injection in larvae resulted in much increased mortality.

Conclusion: According to the findings of this study, insects injected with LAgNPs died at a substantially higher rate than insects injected with latex serum. LAgNPs were efficient against *S. littoralis* larvae and can thus be utilized to specifically control the pest.

Introduction

The cotton leafworm, *Spodoptera littoralis*, is one of the most serious agricultural pests in the Middle East, tropical and subtropical Africa [1]. It is a very harmful pest of many crops including cotton, alfalfa, peanut, potato, lettuce, celery, pepper and tomato. Consequently, several researchers have looked into the biology of *S. littoralis* and the impact of various host plants on their development and reproduction [2,3]. Several chemical studies have been conducted to manage *S. littoralis*, but these synthetic pesticides have resulted in a slew of issues, including pollution, pesticide resistance, and negative impacts on beneficial species like honeybees and birds. As a result, we are applying natural repellents and antifeedants against *S. littoralis* larvae, such as latex from *Calotropis procera*.

C. procera (Aiton) (family Apocynaceae), also known as Sodom apple, usher, and milk weed, is found in tropical and subtropical Africa, Asia, and America. It is a hardy, pubescent, evergreen, erect, compact shrub that grows to a height of 4.5 meters, with a simple, rarely branching stem that is woody at the base and covered with cracks. When the leaves, stem, and flower of this plant are cut or damaged, they can generate a huge amount of latex [4-6].

Numerous plants contain a variety of physiologically active chemicals [7]. The existence of various types of diterpenes, triterpenes, cardiac glycosides, alkaloids, glucosinolates, tannins, saponins, phenolics, and flavonoids has been linked to biological activity in many situations [8]. *C. procera* latex possesses a variety of biological properties, including antibacterial activity [9], anticancer [10], anti-inflammatory [11-13], anthelmintic [14,15], antidiarrheal [16], analgesic [17], antipyretic [18], insecticidal [19-22], antidiabetic [17], hepato-protective [23], schizonticidal [15,17] and antioxidant activities [6,24]. The latex of this plant includes a number of poisonous chemicals called cardiac glycosides (CGs), which act as inhibitors of Na, K-ATPase, and boost intracellular sodium levels. The increase in cardiac contractility is related to an increase in calcium ions in cardiac myocytes [25,26].

Different latex proteins appear to be involved in insect defense strategies [27]. Latex has also been proved to repel insects [28].

In recent years, the world has shifted swiftly to nanotechnology to give answers in a variety of industries, including cosmetics, agriculture, pharmaceuticals, food, and paint. Nanoparticles have an atomic or molecular size of fewer than 100 nanometers [29]. Nanoparticles have attracted a lot of attention because of their unique and intriguing qualities compared to their rivals in terms of applicability. Plant infections, weeds, and insect pests have all been found to be susceptible to nanoparticles. Insecticides and

insect repellents have also included them in their compositions. Fortunately, unlike traditional chemical insecticides, nanoparticles offer no health risks to the environment or the general population [30,31]. However, the exact mechanics of nanoparticles have yet to be fully understood, necessitating in-depth examination or experiments of their interaction with biological systems prior to the development of edible goods.

The purpose of this study is to investigate the potential of utilizing latex serum and latex-synthesized nanoparticles of *C. procera* to manage *S. littoralis*, which is a significant agricultural pest. The study aims to assess the repellent and antifeedant properties of these substances against fourth instar larvae of *S. littoralis*. Additionally, the study contributes to the broader field of nanotechnology in agriculture by exploring the use of nanoparticles as a targeted, safe, and effective means of pest management.

Methods

Insect rearing

The establishment of a stock colony of *S. littoralis* was begun using eggs sourced from the Research Division of the Cotton Leafworm at the Plant Protection Research Institute in Assiut, Egypt. Prior to commencing the studies, the larvae were cultivated in the insectaries located within the Zoology Department of the Faculty of Science at Assiut University. This rearing process spanned 10 generations and involved the utilization of castor-bean leaves, which belong to the Euphorbiaceae family. The adult participants were provided with a sucrose solution of a concentration of 10%. The insects were kept under controlled conditions, with a temperature of $25 \pm 2^\circ\text{C}$, relative humidity of $65 \pm 5\%$, and a photo period of 16 hours of light followed by 8 hours of darkness. An oleander branch, namely from the species *Nerium oleander* L. (Apocynaceae), was introduced inside the cage to serve as a site for oviposition. The egg masses were collected daily and thereafter stored in 90 mL plastic cups at 25°C till the hatching process occurred.

Sampling and fractionation of crude latex

The latex of *C. procera* plant was obtained by tapping method and fractionated into three layers: top layer containing natural rubber; clear middle layer of serum, and the bottom of the lipids [32].

Synthesis and characterization of latex silver nanoparticles (LAg-NPs)

The synthesis and characterization of the synthesized silver nanoparticles has been described before [5].

Antifeeding assay

The antifeeding properties of the latex serum and its nanoparticles (LAG-NPs) were evaluated against the 4th instar larvae of *S. littoralis* using leaf-dip technique. For latex serum, one hundred and eighty larvae were starved overnight, divided into 3 groups each contained 60 larvae, two groups for the treatments one group received latex serum (5, 2.5, 1.25, 0.75 mL) and second group for control. Fresh castor bean leaves of almost equal size were dipped in each treatment and control for 20 seconds before being placed in the shade to dry. The dried leaves were placed in plastic cups one by one. For three days, ten larvae were placed in each cup and allowed to feed on the treated and untreated leaves. For each treatment, six duplicates were carried out. For LAG-NPs, a total of 120 larvae were subjected to an overnight period of starvation. These larvae were then separated into three groups, with each group consisting of 40 larvae. Two of these groups were assigned to experimental treatments, while the remaining group served as the control. Two concentrations of LAG-NPs were used. The weight of each larva and leaf was measured both before and after a 24-hour eating period. The quantification of food consumption and larval mass was determined.

The antifeedant index (AFI) was calculated according to [33].

$$\%AFI = [(C-T) / (C + T)] \times 100$$

Where: C: the amount of food consumed (leaves) in the control

T: the amount of food consumed (leaves) in the treatment

Starvation percentage assay

This bioassay was designed to check the starvation percentage of the 4th instar larvae of *S. littoralis* fed on latex serum and its nanoparticles using leaf-dip technique. For latex serum, a total of 240 larvae were subjected to an overnight period of starvation. These larvae were then separated into four groups, with each group consisting of 60 larvae. Two of the groups were assigned for experimental treatments, one group served as a control, and the remaining group consisted of starved larvae. Fresh castor bean leaves of equal size were dipped in each treatment and control for 20 seconds before being placed in the shade to dry. Each dried leaf was placed in its own plastic cup. The treated and untreated leaves were fed to ten larvae in each cup for 24 hours, whereas the starved larvae went without food for the same amount of time. For each treatment, six duplicates were carried out. All larvae were weighed before to treatment.

For LAG-NPs, one hundred and sixty larvae were starved overnight and were separated into four groups, each with 40 larvae: two for the treatments, one for control, and one for starvation. The latex nanoparticles

were utilized in two concentrations. The larvae were subsequently subjected to reweighing, and the starving percentages of the larvae under investigation were determined using the equation provided by reference [1].

$$\text{Starvation (\%)} = C - E/C - S \times 100$$

Where:

C = Mean weight gain of untreated larvae after 24 h

E = Mean weight gain of larvae treated with latex serum after 24 h

S = Mean weight gain of untreated starved larvae after 24 h.

Topical and injection application of *S. littoralis* with latex serum or latex nanoparticles

All treatments for each experiment were performed on 4th instar larvae from the same batch. Four batches of ten fourth instar larvae were obtained, each containing 40 individuals. On the dorsal side of the abdomen, the larvae were given a single dosage of latex serum or latex nanoparticles (LAG-NPs) (5 % or 10 %). In the case of latex serum treatment, control insects were given a single dosage of 5 µL, 10 µL, or 20 µL of silver nitrate, whereas untreated insects were used as controls. Another experiment involved injecting 5 µL or 10 µL of latex serum or LAG-NPs (5% or 10%) between the second and third abdominal sternites. As a positive control, control larvae were given 5 µl or 10 µl of silver nitrate, whereas untreated larvae were used as a negative control.

Statistical analysis

Percent corrected control was calculated using Abbott's formula [34]. The results were analyzed by ANOVA and Least Significant Difference test (LSD) at P < 0.05 using SPSS.

Results

Synthesis and Characterization of latex silver nanoparticles (LAG-NPs)

The latex silver nanoparticles (LAG-NPs) were synthesized.

UV-vis spectral analysis

The synthesis of LAG-NPs was validated by UV-vis spectroscopic investigations of the colored solution, which revealed distinct surface plasmon resonance bands with a peak centered at roughly 400 nm (Fig. 1). Silver ion reduction was indicated by this color.

Repellent and antifeedant activity of latex serum

In laboratory tests, the repellent and antifeedant activity of *C. procera* latex serum was tested against *S. littoralis* 4th instar larvae during a 24-hour period (Tables 1 and 2). At high concentrations, *C. procera* latex serum had substantial repellent activity ranging from

99.3 to 68.1 %, with starvation percentages of 94.1% and 78.43%, but lower concentrations, repellency ranged from 8.8% to 37.9%, while starvation was observed at 94.1% and 78.43% (corresponding to 21.56% and 47.0%, respectively). In the present study, Data in table 2 indicated that the mean % repellency and starvation was increased with increasing in concentration of latex serum (Fig 2).




Figure 1: Absorption spectrum solution of silver nanoparticles synthesized by using *C. procera* latex serum after 1 h.

Repellent and antifeedant properties of latex (LAG-NPs)

After 24 hours of feeding on treated castor leaves, the antifeedant activity of the 2 % and 5 % concentrations of LAG-NPs against 4th instar *S. littoralis* larvae is shown in Table 3. Both concentrations had antifeedant characteristics against 4th instar larvae, according to the data. The antifeedant impact of a 5 % concentration was more strong than that of a 2 % concentration, with 5 % having four times the antifeedant activity of 2 %. The starving percentage of the 4th instar larvae treated with the aforementioned concentrations is shown in table (4). The starvation percentage of larvae treated with a 5 % concentration was 17.75 %, whereas it was 9.66 % with a 2 % concentration. As a result, the greater concentration proved to be more effective than the lower one as an antifeedant. (Fig. 3).

Latex serum Conc.(ml)	Antifeedant index (%)			Mean*
	Days post-treatments			
	1 st day	2 nd day	3 rd day	
5	100.0 ± 0.00 a	97.90 ± 0.69 a	100.0 ± 0.00 a	99.3
2.5	76.92 ± 0.89 b	61.42 ± 1.01 b	65.76 ± 1.20 b	68.6
1.25	39.39 ± 0.49 c	30.49 ± 0.86 c	43.75 ± 1.24 c	37.9
0.75	6.350 ± 0.19 d	6.370 ± 0.20 d	13.58 ± 0.73 d	8.80
LSD	0.738	1.075	1.328	
P	0.0000	0.0000	0.0000	
df	3,20	3,20	3,20	

Table 1: Antifeedant activity of latex serum against 4th instar larvae of *S. littoralis* fed on castor leaves treated with (*C. procera*) different latex concentrations. Data are expressed as mean ± SE (n= 6). * General mean of each treatment at different time intervals. Data are statistically analyzed by one-way ANOVA. Means within each column followed by different letters are significantly different based on LSD (P < 0.05).

The effects of topical application of latex serum or its nanoparticles (LAGNPs)

Topical application of different concentrations (5 µl, 10 µl or 20 µl) of latex serum or latex synthesized nanoparticles (LAGNPs) (5 % or 10%) on 4th instar larvae

of *S. littoralis* had no significant effect in mortality in comparison to control insects.

Treatments Conc.	Average weight at zero time (mg/larva)	Average weight after 24 h (mg/larva)	Difference* (mg/larva)	Starvation (%)
5	43	43	0	94.1%
2.5	46	54	+ 8	78.43%
1.25	33	57	+ 24	47.0%
0.75	40	77	+ 37	21.56%
Control larvae	43	91	+ 48	-----
Starved larvae	42.0	39.0	- 3	-----

*Average weight after 24 h – Average weight at zero time.

Table 2: Starvation percentage (%) of the 4th larval instar of *S. littoralis* as affected by *C. procera* latex serum.




Figure 2: Antifeedant and starvation percentage (%) activity of *C. procera* latex serum against 4th instar larvae of *S. littoralis* fed on castor leaves treated with different latex concentrations.

Latex nanoparticles Conc.(per mg)	Antifeedant index (%)			Mean*
	Days post-treatments			
	1 st day	2 nd day	3 rd day	
2	2.920 ± 0.19 b	5.190 ± 0.38 b	9.832 ± 0.41 b	6.00
5	28.50 ± 0.87 a	28.50 ± 1.24 a	27.33 ± 0.73 a	28.5
LSD	0.894	1.306	0.845	
P	0.0000	0.0000	0.0000	
df	1,6	1,6	1,6	

Table 3: Antifeedant activity of LAG-NPs (*C. procera*) against 4th instar larvae of *S. littoralis* fed on castor leaves treated with different concentrations. Data are expressed as mean ± SE (n= 4).

* General mean of each treatment at different time intervals. Data are statistically analyzed by one-way ANOVA. Means within each column followed by different letters are significantly different based on LSD (P < 0.05).

Treatments Conc.	Average weight at zero time (mg/larva)	Average weight after 24h (mg/larva)	Difference* (mg/larva)	Starvation (%)
2 %	54.33	86.26	+ 31.93	9.66 %
5 %	47	75.52	+ 28.52	17.75 %
Control larvae	54	90	+ 36	-----
Starved larvae	40.13	34	- 6.13	-----

*Average weight after 24h – Average weight at zero time.

Table 4: Starvation percentage (%) of the 4th larval instar of *S. littoralis* as affected by LAG-NPs (*C. procera*).

The effects of injection of latex serum or its nanoparticles (LAGNPs)

The efficacy of *C. procera* latex serum and LAGNPs produced from this latex serum against *S. littoralis* 4th

instar larvae were tested. Figure (4) shows the findings of a serum larvicidal bioassay, while Figure 5 A and B shows the results of latex-synthesized AgNPs. A single dosage of 5 μ l of latex serum injected into 4th instar larvae was unsuccessful ($P = 0.267$), but a single dose of 10 μ l of latex serum injected into 4th instar larvae resulted in a minor influence ($P = 0.077$) in mortality percentage on day 2 following treatment when compared to control insects (Fig. 4). Furthermore, on day 3 following treatment, there was a significant difference ($P = 0.002$) in mortality percentage when compared to control insects (Fig. 4).




Figure 3: Antifeedant and starvation percentage (%) activity of *C. procera* latex nanoparticles against 4th instar larvae of *S. littoralis* fed on castor leaves treated with different concentrations.





Figure 4: Mortality percentage (mean $\pm$ SE) of *S. littoralis* larvae caused by different doses of latex serum. Bars with different letters were significantly different ($P < 0.05$) by LSD after ANOVA.

A



B




Figure 5: Mortality percentage (mean $\pm$ SE) of *S. littoralis* larvae caused by different doses of 5% LAgNPs (A) and 10% LAgNPs (B). Bars with different letters were significantly different ($P < 0.05$) by LSD after ANOVA.

Injection of a single dosage of 5% (5 μ l, or 10 μ l) or 10% (5 μ l, or 10 μ l) of LAgNPs, on the other hand, had a significant influence on mortality when compared to controls (Fig. 5 A and B). In general, the mortality rate has risen steadily over time. With rising LAgNPs concentrations, the percentage mortality rose, showing a direct link between dosage and percent mortality (Fig. 5 A and B). Furthermore, on days 1 and 2, there was no statistically significant difference observed in the percentage of mortality between the untreated control (negative control) and the silver nitrate-treated control (positive control), but there was a significant difference ($P = 0.009$) on day 3 of treatment (Fig. 5 A and B).

Discussion

Insects injected with LAgNPs had much greater mortality than insects injected with latex serum, according to the findings of this study. This might be due to the larvae's improved protection system against injected latex serum due to more efficient detoxifying. This, on the other hand, is supported by the larvicidal behavior of AgNPs. The high surface area-to-volume

ratio of AgNPs confers a wide range of biocidal and catalytic properties. In latex-mediated nanosynthesis, encapsulating latex metabolites on the surface of AgNPs improves their larvicidal activity in addition to providing stability [35].

C. procera is a well-known medicinal plant, with several physiologically active chemicals in its latex [5] used Transmission Electron Microscopy (TEM) analysis, laser diffraction particle size analyzer, X-ray diffraction (XRD) analysis, and FTIR spectrophotometer to characterize latex nanoparticles.

The repellent and antifeedant activity of *C. procera* latex serum has been recorded in *C. procera* latex with high concentrations of active compounds such as Terpenes, phenolic compounds, and cardiac glycosides, and these compounds play a role in cotton leaf worm repellent and antifeedant [6].

Plants have a multitude of compounds that each have their own biological action. *C. procera* latex was discovered to contain a variety of chemicals, including cardenolides, proteolytic enzymes, alkaloids, and carbohydrates, according to many studies [35].

C. procera latex has also been found to work as repellents and oviposition deterrents by a number of studies [36]. Moreover, [37] showed that the latex of *C. procera* possessed high repellent activity against larvae of *Culex quinquefasciatus*.

Compared to control insects, topical administration of latex serum or its nanoparticles (LAgNPs) showed no significant influence on mortality. This might be due to improved peripheral latex or LAgNP protection mechanisms, such as more effective metabolic detoxification, faster excretion rates, and reduced integument penetration rates. In contrast to the current findings, *C. procera* latex had insecticidal action against *Musca domestica* 3rd instar larvae at a topically dosage of 3 µL (5% of the latex) [35].

In general, the 4th instar larvae of *S. littoralis* showed a greater sensitivity to LAgNP injection than to latex serum injection. In this respect, [38] reported that latex materials from different plants (*Euphorbia milii*, *E. hirta*, *Ficus racemosa* and *Jatropha curcas*) were less toxic than the synthesized AgNPs to the 2nd and 4th instar larvae of *Aedes aegypti* and *Anopheles stephensi*. In addition, [39,40] reported larvicidal activity of silver nanoparticles synthesized by *Pergularia daemia* latex against *Aedes aegypti* and *Anopheles stephensi*.

The latex serum utilized in this investigation had significant levels of terpenes, phenolic compounds, and cardiac glycoside, as previously reported [6], so a preliminary conclusion could be drawn that there is an interaction between these compounds in AgNPs formation. Furthermore, *C. procera* Latex serum includes "cardiac glycoside," one of the most bioactive substances, which predominantly affects the

cardiovascular, neurologic, and gastrointestinal systems [41].

Cellular depolarization and the loss of the negative membrane potential essential for proper cell activity would result if these ion gradients were lost. More research on the mechanism of silver nanoparticles' larvicidal effect is needed. Additionally, more studies should evaluate nanoparticles that are naturally occurring and easily available for IPM.

Food security concerns may benefit from nanotechnology, such as targeted pesticide delivery and control of pestiferous insects that destroy crops and their products in the field. The current study demonstrated that the insects injected with LAgNPs died at a significantly greater rate than insects injected with latex serum, according to the findings of this study. LAgNPs were effective against *S. littoralis* larvae and can thus be used to control the pest specifically.

Acknowledgements

The authors grateful to Assiut University for funding and technical support and to Asmaa Metwaly, Faculty of Science, Assiut University for contributing the experiment tools and insect rearing. The authors declare that we have no conflict of interest.

Funding

No special funding was obtained for this study.

Author Contributions

Amer Tawfik: contributed to conceptualization, Nadia Mohamed and Ahmed Ali: contributed to methodology, Ahmed Ali and Mady Ismail: contributed to statistical analysis writing, Amer Tawfik and Ahmed Ali: contributed to writing, review and editing, Wael. M. Abdel-Mageed and Ahmed Shoreit: contributed to project administration. All authors have read and agreed to the published version of the manuscript.

Conflicts of interest

The authors declare no conflict of interest.

References

1. Abdel-Mageed MI, Shaaban AM, Zidan ZH. The effectiveness of Dursban, Du-ter Brestan and their combination against the cotton leafworm *Spodoptera littoralis* (Boisd). Bulletin of the Entomological Society of Egypt, Economic Series, (1975); 9:283-292.
2. Mohamed H. Comparative study of host plants on growth, development and fecundity of the cotton leafworm *Spodoptera littoralis*, (Boisduval) Noctuidae: Lepidoptera. Journal of the Egyptian German Society of Zoology, (2003); 42(E): 167-184.
3. Adham FK, Rashad EM, Shoukry IF, Nasr EE. Host plants shifting affects the biology and biochemistry of *Spodoptera littoralis* (boisd.)(lepidoptera: noctuidae). Egyptian

- Academic Journal of Biological Sciences A, Entomology, (2009); 2(1): 63-71.
4. Tour N, Talele G. Anti-inflammatory and gastromucosal protective effects of *Calotropis procera* (Asclepiadaceae) stem bark. *Journal of Natural Medicines*, (2011); 65:598-605.
 5. Mohamed NH, Ismail MA, Abdel-Mageed WM, Shoreit AAM. Antimicrobial activity of latex silver nanoparticles using *Calotropis procera*. *Asian Pacific Journal of Tropical Biomedicine*, (2014); 4(11): 876-883.
 6. Sayed AE-DH, Mohamed NH, Ismail MA, Abdel-Mageed WM, Shoreit AA. Antioxidant and antiapoptotic activities of *Calotropis procera* latex on Catfish (*Clarias gariepinus*) exposed to toxic 4-nonylphenol. *Ecotoxicology and environmental safety*, (2016); 128:189-194.
 7. ÖKSÜZ S, Ulubelen A, Barla A, VOELTER W. Terpenoids and aromatic compounds from *Euphorbia heteradana*. *Turkish Journal of Chemistry*, (2002); 26(4): 457-464.
 8. Lv D, Liu X, Dong Y, Yan Z, Zhang X, et al. Comparison of gut bacterial communities of fall armyworm (*Spodoptera frugiperda*) reared on different host plants. *International Journal of Molecular Sciences*, (2021); 22(20): 11266.
 9. Manoorkar V, Mandge S, Gachande B. Antifungal activity of leaf and latex extracts of *Calotropis procera* (Ait.) against dominant seed-borne storage fungi of some oil seeds. *Bioscience discovery*, (2015); 6(1): 22-26.
 10. Shaker KH, Morsy N, Zinecker H, Imhoff JF, Schneider B. Secondary metabolites from *Calotropis procera* (Aiton). *Phytochemistry Letters*, (2010); 3(4): 212-216.
 11. Arya S, Kumar VL. Antiinflammatory efficacy of extracts of latex of *Calotropis procera* against different mediators of inflammation. *Mediators of inflammation*, (2005); 4(4): 228-232.
 12. Kumar VL, Roy S. *Calotropis procera* latex extract affords protection against inflammation and oxidative stress in Freund's complete adjuvant-induced monoarthritis in rats. *Mediators of inflammation*, (2007); 1-7.
 13. Abdel-Mageed WM, Mohamed NH, Liu M, El-Gamal AA, Basudan OA, et al. Lipoxygenase inhibitors from the latex of *Calotropis procera*. *Archives of Pharmacol Research*, (2016); 1-9.
 14. Shivkar Y, Kumar V. Anthelmintic activity of latex of *Calotropis procera*. *Pharmaceutical biology*, (2003); 41(4): 263-265.
 15. Murti Y, Sharma S, Mishra P. In vitro anthelmintic activity of *Calotropis procera* (AIT.) R. BR. Leaves. *Asian Journal of Pharmaceutical and Clinical Research*, (2015); 8(6).
 16. Das RK, Sharma P, Nahar P, Bora U. Synthesis of gold nanoparticles using aqueous extract of *Calotropis procera* latex. *Materials Letters*, (2011); 65(4): 610-613.
 17. Quazi S, Mathur K, Arora S, Wing P. *Calotropis procera*: An overview of its phytochemistry and pharmacology. *Indian Journal of Drugs*, (2013); 1(2): 63-69.
 18. Larhsini M, Markouk M, Jaouhari J, Bekkouche K, Lazrek H, et al. The antipyretic activity of some Moroccan medicinal plants. *Phytotherapy Research*, (2002); 16(S1): 97-98.
 19. Moursy LE. Insecticidal activity of *Calotropis procera* extracts of the flesh fly, *Sarcophaga haemorrhoidalis* fallen. *Journal of the Egyptian Society of Parasitology*, (1997); 27(2): 505-514.
 20. Singh R, Mittal P, Dhiman R. Laboratory study on larvicidal properties of leaf extract of *Calotropis procera* (Family-Asclepiadaceae) against mosquito larvae. *Journal of Communicable Diseases*, (2005); 37(2): 109.
 21. Ramos MV, Bandeira GdP, Freitas CDTd, Nogueira NAP, Alencar NMN, et al. Latex constituents from *Calotropis procera* (R. Br.) display toxicity upon egg hatching and larvae of *Aedes aegypti* (Linn.). *Memórias do Instituto Oswaldo Cruz*, (2006); 101:503-510.
 22. Shahi M, Hanafi-Bojd AA, Iranshahi M, Vatandoost H, Hanafi-Bojd MY. Larvicidal efficacy of latex and extract of *Calotropis procera* (Gentianales: Asclepiadaceae) against *Culex quinquefasciatus* and *Anopheles stephensi* (Diptera: Culicidae). *Journal of Vector Borne Diseases*, (2010); 47(3): 185-188.
 23. Setty SR, Quereshi AA, Swamy AV, Patil T, Prakash T, et al. Hepatoprotective activity of *Calotropis procera* flowers against paracetamol-induced hepatic injury in rats. *Fitoterapia*, (2007); 78(7-8): 451-454.
 24. Chaudhary P, de Araújo Viana C, Ramos MV, Kumar VL. Antiedematogenic and antioxidant properties of high molecular weight protein sub-fraction of *Calotropis procera* latex in rat. *Journal of Basic and Clinical Pharmacology*, (2015); 6(2): 69.
 25. Juncker T, Schumacher M, Dicato M, Diederich M. UNBS1450 from *Calotropis procera* as a regulator of signaling pathways involved in proliferation and cell death. *Biochemical Pharmacology*, (2009); 78(1): 1-10.
 26. Cerella C, Muller F, Gaigneaux A, Radogna F, Viry E, et al. Early downregulation of Mcl-1 regulates apoptosis triggered by cardiac glycoside UNBS1450. *Cell death & disease*, (2015); 6(6): e1782-e1782.
 27. Konno K, Hirayama C, Nakamura M, Tateishi K, Tamura Y, et al. Papain protects papaya trees from herbivorous insects: role of cysteine proteases in latex. *The Plant Journal*, (2004); 37(3): 370-378.
 28. Singhi M, Joshi V, Sharma R, Sharma K. Ovipositioning Behaviour of *Aedes aegypti* in Different Concentrations of Latex of *Calotropis Procera*: Studies on Refractory Behaviour and its Sustenance across Gonotrophic Cycles. (2004).
 29. Gutiérrez-Ramírez JA, Betancourt-Galindo R, Aguirre-Urbe LA, Cerna-Chávez E, Sandoval-Rangel A, et al. Insecticidal effect of zinc oxide and titanium dioxide nanoparticles against *Bactericera cockerelli* Sulc. (Hemiptera: Trioziidae) on tomato *Solanum lycopersicum*. *Agronomy*, (2021); 11(8): 1460.
 30. Barik T, Sahu B, Swain V. Nanosilica—from medicine to pest control. *Parasitology Research*, (2008); 103:253-258.
 31. Thabet AF, Boraie HA, Galal OA, El-Samahy MF, Mousa KM, et al. Silica nanoparticles as pesticide against insects of different feeding types and their non-target attraction of predators. *Scientific Reports*, (2021); 11(1): 1-13.
 32. Kang H, Kang MY, Han K-H. Identification of natural rubber and characterization of rubber biosynthetic activity in fig tree. *Plant physiology*, (2000); 123(3): 1133-1142.
 33. Sadek MM. Antifeedant and toxic activity of *Ahadotoda vasica* leaf extract against *Spodoptera littoralis* (Lep., Noctuidae). *Journal of applied entomology*, (2003); 127(7): 396-404.
 34. Abbott WS. A method of computing the effectiveness of an insecticide. *Journal of economic Entomology*, (1925); 18(2): 265-267.
 35. Morsy TA, Rahem M, Allam K. Control of *Musca domestica* third instar larvae by the latex of *Calotropis procera* (Family: Asclepiadaceae). *Journal of the Egyptian Society of Parasitology*, (2001); 31(1): 107-110.
 36. Kabir M, Iqbal MZ, Farooqi Z, Shafiq M. Vegetation pattern and soil characteristics of the polluted industrial area of Karachi. *Pakistan Journal of Botany*, (2010); 42(1): 661-678.
 37. Tahir HM, Ishaq T, Muhammad KM, Khan SY, Ahmed K. Potential use of *Calotropis procera* (milk weed) to control *Culex quinquefasciatus* (Diptera: culicidae). *Pakistan Journal of Zoology*, (2013); 45(3).
 38. Borase H, Patil C, Salunkhe R, Narkhede C, Suryawanshi R, et al. Mosquito larvicidal and silver nanoparticles synthesis potential of plant latex. *Journal of Entomological and Acarological Research*, (2014); 46(2): 59-65.
 39. Patil CD, Borase HP, Patil SV, Salunkhe RB, Salunkhe BK. Larvicidal activity of silver nanoparticles synthesized using *Pergularia daemia* plant latex against *Aedes aegypti* and *Anopheles stephensi* and nontarget fish *Poecilia reticulata*. *Parasitology Research*, (2012); 111:555-562.
 40. Patil CD, Patil SV, Borase HP, Salunkhe BK, Salunkhe RB. Larvicidal activity of silver nanoparticles synthesized using

Plumeria rubra plant latex against Aedes aegypti and Anopheles stephensi. Parasitology Research, (2012); 110:1815-1822.

41. Wang Z-N, Wang M-Y, Mei W-L, Han Z, Dai H-F. A new cytotoxic pregnanone from Calotropis gigantea. Molecules, (2008); 13(12): 3033-3039.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License. To read the copy of this license please visit: <https://creativecommons.org/licenses/by-nc/4.0/>