



Full Length Research Article

Advancements in Life Sciences – International Quarterly Journal of Biological Sciences

ARTICLE INFO

Open Access



Date Received:

04/05/2023;

Date Revised:

09/06/2023;

Date Published Online:

30/09/2023;

Authors' Affiliation:

Liaquat University of Medical and Health Sciences, Jamshoro, Sindh - Pakistan

*Corresponding Author:

Fahmida Gul

Email:

fahmida.gul@lumhs.edu.pk

How to Cite:

Gul F, Memon S, Ujjan I, Goswami P, Bhatti KA (2023). The effects of 4 β -Hydroxy withanolide E extracted from *Physalis Peruviana* on Complete Blood Count of Dimethylbenz(a)anthracene-induced Breast Cancer in Albino Rats. Adv. Life Sci. 10(3): 434-438.

Keywords:

Physalis Peruvian; Breast cancer; complete blood counts; Dimethylbenz(a)anthracene

The effects of 4 β -hydroxy withanolide E extracted from *Physalis Peruviana* on Complete Blood Count of Dimethylbenz(a)anthracene-induced Breast Cancer in Albino Rats

Fahmida Gul*, Samreen Memon, Ikramuddin Ujjan, Pushpa Goswami, Kanwal Abbas Bhatti

Abstract

Background: Several studies reveal changes in blood parameters in patients with cancer. Complete blood count is a good predictor for many cancer diagnoses and prognoses. They are also helpful in assessing drug efficacy and toxicity. Cancers may be treated with bioactive and antioxidant compounds derived from plants. The effects of *Physalis peruviana* on the complete blood count of dimethylbenz(a)anthracene induced are still not well identified.

Methods: This study was conducted at Liaquat University of Medical and Health Sciences, Jamshoro, for six months on 60 female albino rats divided in 05 groups. The effects of *Physalis peruviana* fruits and tamoxifen on complete blood count in dimethylbenz(a)anthracene induced breast cancer were assessed. Blood samples were collected at weeks 1, 5, 7, and 10 for hematological changes before, during, and after treatment.

Result: Rats of group E (Tumor-induced albino rats treated with the extracts of *Physalis Peruviana* and tamoxifen) gave the best results compared to other groups. In this group, increases in weight along with an improvement in hematological parameters were observed when compared with others.

Conclusion: The extract from *Physalis Peruviana* helps in the reversal of pancytopenia with anemia and thrombocytopenia caused by tumor-inducing drugs.



Introduction

In women, breast cancer is the most common form of malignancy. Breast cancer is the second leading cancer-related cause of death for women worldwide. Approximately one in nine women suffers from breast cancer. With an incidence rate of 50/100,000 in Pakistan, 83,000 cases registered with a death rate of over 40,000 [1]. There are many therapeutic methods including surgical removal, radiation and chemotherapy. This troubling condition challenges healthcare workers to search for a new cure and remedy. Among many others, the oldest anticancer remedies from extracts of several plants are drawing attention due to potential side effects and resistance of current therapies [2].

Physalis peruviana belongs to the Solanaceae family and the *Physalis* genus known as golden berry and Cape gooseberry [3]. The anti-proliferative effects of *Physalis peruviana* leaves, stems, and whole plants on various carcinomas including colon, chronic myeloid leukemia, and breast cancer are well recognized. Its bioactive compound, 4β-Hydroxy withanolide E generates reactive oxygen species (ROS) inducing apoptosis of cancer cells. The accumulation of this bioactive compound in tumor cells contributes to regulating Nrf2 and superoxide dismutase protein expression. The presence of catalase leads to decrease in tumor size [4]. The blood is a vital connective tissue; a basic medium of exchange for oxygen, carbon dioxide, nutrients, and metabolites essential for life [5]. Several studies report various side effects on blood parameters mostly red blood cells due to cancer therapy [6]. This study is intended to determine various hematological parameters of DMBA-induced breast cancer and to assess the effects of *Physalis peruviana* extract 4β-Hydroxy withanolide E on various hematological parameters.

Methods

Study place and duration

This study was conducted after institutional ethical approval for six months at the Liaquat University of Medical and Health Sciences, Jamshoro in Pakistan in collaboration with Sindh Agricultural University Tando Jam Animal House and the Medical Research Center Jamshoro, Pakistan.

Animals: For this study, 60 adult female Wistar albino rats were purchased from the National Institute of Health, Karachi. Healthy female rats weighing between 165gms to 175gms were included in this study.

Experimental design: Animals were handled according to protocol of animal's care provided by guidelines of National Institute of Health Pakistan (NIH). Stainless

steel cages were used for housing animals with 20 mm thick rice husk bedding in each cage; well ventilated and had a 12-hour light-dark cycle alternatively with temperature 22 ± 2 degrees Celsius, and relative humidity of $55\pm 10\%$ [7]. The environment of the cages and the surroundings were kept properly hygienic and adequately ventilated. Each cage contained four animals. The food and water were accessible to the animals to *ad libitum*. The standard chow was used as feed and tap water to drink. Animals were divided in five equal groups, each group comprised of 12 animals, housed in separate cages. All animals were weighed and a blood sample was taken before, during and after experiment [8].

Animal Groups and Treatments:

Group A: PC (Positive control) – Control experiment with normal diet and water.

Group B: NC (Negative control) – DMBA induced rats only not treated.

Group C: TR1– DMBA Induced rats were treated with tamoxifen.

Group D: TR2– DMBA induced rats were treated with 4 β-Hydroxy withanolide E

Group E: TR3– DMBA induced rats were treated with tamoxifen and 4 β-Hydroxy withanolide E.

Plant Material and Extract Preparation: *Physalis peruviana* was obtained from Botanical store Karachi, Pakistan, certified by the Department of Botany of Sindh University Jamshoro. One hundred grams of *Physalis peruviana* was diluted in 1 liter boiling water. Filtered extract under the filter paper and dried for 3 hours and stored at -21°C until use.

Tumor Induction: DMBA (Dimethylbenz(a)Anthracene) is a yellowish green, odorless chemical derivative of phenanthrene; an immunosuppressant and a potent laboratory carcinogen purchased from Sigma Aldrich.

Soy oil: purchased from the Departmental store.

A single dose of DMBA (20 mg) in soy oil (5 ml) given by intra-gastric gavage is sufficient to induce tumor in the mammary gland. Each rat was weighed and examined for tumor in the mammary gland by gross inspection, palpation weekly. The tumor was confirmed by biopsy on the 6th week after an administration of DMBA [9].

***Physalis Peruviana* administration:** Extracted dried *Physalis peruviana* 100 mg/kg/week administered with diet.

Tamoxifen administration: Tamoxifen (Sigma-Aldrich) dissolved in soy oil at a concentration of 20 mg/ml by shaking overnight at 37°C . 75 mg tamoxifen/kg body weight via intraperitoneal injection was administered.

Blood sampling: The blood samples were collected before tumor induction and at weeks 1, 5, 7, and 10 after anesthetizing rats with ketamine injection. The blood (1ml) was collected by retro-orbital technique via the orbital sinus in heparinized micro-crit capillary tubes and transferred to the vacutainer blood collection tube [10].

Statistical analysis:

The data were analyzed using SPSS version 26.0. Numerical parameters like weight, hemoglobin, platelets, red blood cells, and white blood cells were presented as mean \pm SD and one-way analysis of Variance (ANOVA) and T-test were applied to compare the means between the groups. P-value < 0.05 was considered as a statistically significant level for all comparisons.

Results

We assessed the difference in body weight measurement (table 1) and hematological parameters; Hemoglobin (Hb), Red Blood Cells (RBCs), White Blood Cells (WBCs) and platelets with Mind-ray hematology analyzer - BC-2300 (table 2).



Figure 1: The administration of *Physalis peruviana* and collection of the blood sample.

Weight measurement

Mean of the weight was compared among the five groups by applying one-way ANOVA, and there was a significant increase in the weight of rats of Group E versus Group B (p-value = 0.0001). Negative Control showed massive weight loss and worsening of all hematological parameters like pancytopenia, anemia and leucopenia. Six rats died during the 10th week because of the severity of the illness.

Hematological parameters

By applying the ANOVA test, significant decrease in RBC and hemoglobin in both B and C Groups (P-value = 0.0001) was observed. There was a significant increase in RBCs and hemoglobin in Group E. WBC and platelets have increased to normal levels after treatment with *Physalis peruviana*. We found a significant decrease in WBC and Platelets in Group B and Group D (P-value = 0.0001). A significant increase in WBC and platelets in

Group E (P-value = 0.0001) was also observed after 10 weeks. There were no major variations between Group A and Group E. After treatment, Group E revealed gross improvement in weight and all hematological parameters.

Abbreviations: PC, positive control (Normal); NC, negative control (Induced with DMBA, no treatment) – Control experiment; TR1, DMBA induced rats treated with tamoxifen; TR2, DMBA induced with 4 β -hydroxy withanolideE; TR3, DMBA induced with tamoxifen and 4 β -hydroxy withanolide E

Discussion

The phytochemicals found in leaves, fruits, and other parts of several plants have therapeutic potential for treating various diseases including cancers [11]. The incidence of breast cancer is increasing in developing countries [12]. Several factors contribute to cancer development, including inflammation [13]. In breast tissue, DMBA causes DNA damage and promotes cell proliferation [14]. Vitamin C & E, polyphenols, carotenoid compounds, lutein, and lycopene are antioxidants present in many plant extracts, helps in tumor suppression [15].

The complete blood count reflects the response of cellular immunity in a patient with cancer. [16]. In this study, after induction with DMBA (carcinogen), blood sampling at weeks 1, 5, 7, and 10 revealed drastic variations among Group A (Positive Control) and Group B (Negative control (Induced with DMBA no treatment)). We found significant improvement in all parameters of Group E (DMBA induced rats treated with tamoxifen and 4 β -hydroxy withanolideE) compared to Group B (Negative control (Induced with DMBA no treatment)). These observations are in line with Akuru [17] who reported indices of DMBA administered albino rats, given aqueous concentrate of different plants. For RBC and Hb concentrations, there was a significant increase (p<0.05) in all the treated groups when contrasted with the control and DMBA groups (p<0.05) [17].

In our results, there was a significant decrease in RBC and hemoglobin in all four groups induced by DMBA compared to the Group A. This contrasts with Akuru et al, who showed contrary results and reported that there was an additional increase in the RBC level in all groups when compared with the control and DMBA untreated group, however not significant (p<0.05) [17]. This study also suggests anemia in cancer with low Hb level and red blood cell count, which may predict the death among cancer patients [18,19]. There is a high incidence of anemia observed in cancer patients, which may be caused by bleeding, nutritional deficiencies, damage to the bone marrow, or tumor infiltration [20].

Weight (grams, mean±SD)	Week	Groups					p-value
		A: PC	B: NC	C: TR1	D: TR2	E: TR3	
	1	190.83±5.14	194.58±23.39	201.25±21.43	198.75±19.67	212.08±12.33	0.054*
	5	209.5±11.5	176.25±26.89	178.75±26.80	175.41±25.71	199.16±12.58	<0.0001*
	7	230.41±14.91	159.16±32.53	191.66±20.92	187.91±21.15	211.66±11.14	<0.0001*
	10	276.25±11.30	261.66±76.46	207.91±17.76	207.08±23.68	233.33±15.71	<0.0001*

* p-value is statistically significant calculated by One Way ANOVA

Table 1: Comparison of weight at week 1,5,7 and 10 in different groups (n=12) of Albino rats

Parameters	Week	Groups					p-value
		A: PC	B: NC	C: TR1	D: TR2	E: TR3	
RBC	1	9.15±0.65	9.50±0.92	9.49±0.76	7.99±0.76	7.79±0.76	0.390
	5	8.45±0.65	8.80±0.92	8.79±0.76	7.29±0.76	7.09±0.76	0.867
	7	9.65±0.61	6.60±1.79	6.55±2.06	7.29±0.76	6.09±0.76	<0.0001*
	10	9.37±0.96	2.95±2.65	4.59±0.80	5.09±0.76	9.05±0.76	<0.0001*
Haemoglobin	1	14.42±1.64	15.17±1.15	14.17±1.15	12.97±1.15	11.67±1.15	<0.0001*
	5	13.42±1.64	14.17±1.15	13.17±1.15	11.97±1.15	10.67±1.15	<0.0001*
	7	15.73±1.82	11.15±1.69	14.48±0.91	13.48±0.91	12.98±0.91	<0.0001*
	10	15.65±1.71	6.30±5.23	44.20±1.51	11.98±0.91	11.48±0.91	<0.0001*
WBC	1	8.01±1.38	7.61±0.98	6.99±1.43	7.21±1.38	6.71±1.38	0.144
	5	7.71±1.38	7.31±0.98	6.69±1.43	6.91±6.91	6.41±1.38	0.154
	7	7.54±0.91	13.66±4.57	14.95±4.04	14.45±4.04	13.95±4.04	<0.0001*
	10	7.01±1.15	2.10±1.88	7.35±2.59	6.95±2.59	13.45±4.04	<0.0001*
Platelet	1	387.83±82.6	370.16±67.63	366.33±23.56	367.83±53.40	693.75±254.5	0.132
	5	387.33±82.6	369.66±7.63	365.83±23.56	367.33±53.40	693.25±254.5	<0.0001*
	7	369.50±62.7	781.66±259.62	666.91±151.29	369.50±62.73	781.66±74.94	<0.0001*
	10	415.33±144.2	655.50±485.37	449.62±161.29	367.50±62.73	779.66±259.62	<0.0001*

* p-value is statistically significant calculated by One Way ANOVA

Table 2: Comparison of Hematological parameters at week 1,5,7 and 10 in different groups(n=12) of Albino rats

Different plants have been shown to possess anti-anemic properties. Zingue et al. observed ethanolic extracts of plants were useful in increasing multiple biochemical markers of DMBA-induced mammary tumors, including MCV, MCHC, MCH, RBC, PCV, and hemoglobin concentrations, compared to DMBA-untreated animals [21].

We observed enhanced WBC representation of lymphocytosis in all groups induced by DMBA. White Blood Cells have increased to normal levels after treatment with *physalis peruviana* and tamoxifen in group E (DMBA induced rats treated with tamoxifen and 4 β-Hydroxywithanolide E) versus Group C (DMBA induced rats treated with tamoxifen) and D (DMBA induced with 4β-hydroxy withanolideE). This was similar to the study conducted by Akuru et al, who revealed the white blood cells of DMBA administered to albino rats given different plant extracts. There was a significant increase ($p < 0.05$) in the white blood cell level of animals treated with 1000mg/kg of a plant when compared with the control and the DMBA-untreated group. The DMBA group had the most reduced white blood cell level(not significant, $p \leq 0.05$) [17]. When the WBC count is below the normal level, infection is a serious risk, whereas higher counts increase the chance of developing invasive breast cancer. This is also consistent with the findings of Chen et al. [22], who found that plant extracts could reduce the levels of WBCs, neutrophils, and lymphocytes in DMBA-induced mammary cancer. In a study, Akuru et al (2019) measured blood platelet concentrations of

albino rats given DMBA treatment and an aqueous extract from different plants. The DMBA-untreated group showed a significant increase ($p \leq 0.05$) when compared to the control group and the DMBA STD group. The DMBA-untreated group had lower values than the various treated groups (although not significantly, $p \leq 0.05$). According to Rochet, Markovic [23] because platelets release a variety of growth factors and cytokines that promote angiogenesis, high platelet counts are associated with poorer prognosis for cancer, which is crucial to breast cancer metastasis [24]. This study reveals that this extract reduced blood platelets and hence angiogenesis, as evidenced by the lower values of platelet count. Similar is reported by Zingue et al [21], that ethanolic extracts of the plant reduced platelet concentration in tumors induced by DMBA compared to the group not exposed to DMBA.

This study concludes *Physalis peruviana* aqueous extract was found to enhance all hematological parameters of DMBA-induced breast cancer in albino rats. It increases the red blood cells and platelet count and stabilizes white blood cells in their function to reduce the anemic burden caused by cancer. It decreases the risk of thrombocytopenia. Compared with different adverse effects and contraindications of traditional drugs, cultural values, efficacy, low cost, and easy availability encourage the use of medicinal herbs. The use of *Physalis peruviana* in breast cancer treatment has significant potential, especially the adjunct to tamoxifen. To evaluate these effects, further research is required, particularly clinical studies.

Competing Interest

The authors declare that there is no conflict of interest.

Author Contributions

Fahmida Gul: Designed the experiment, data collection and analyzes the study and drafting of the manuscript.

Corresponding author

SamreenMemon: Conceived the idea, was the in-charge of data collection, final approval of the manuscript.

Ikramuddin Ujjan; Brought new ideas, helped in technical issues.

PushpaGoswami: Helped in data collection and drafting of manuscript.

Kanwal Abbas: Helped in statistics for data analysis and final drafting of manuscript

References

- Hu C, Hart SN, Gnanaolivu R, Huang H, Lee KY, et al. A Population-Based Study of Genes Previously Implicated in Breast Cancer. *N Engl J Med*, (2021); 384(5): 440-451.
- Waks AG, Winer EP. Breast Cancer Treatment: A Review. *JAMA*, (2019); 321(3): 288-300.
- El-Beltagi HS, Mohamed HI, Safwat G, Gamal M, Megahed BMH. Chemical Composition and Biological Activity of *Physalis peruviana* L. *Gesunde Pflanzen*, (2019); 71113-122.
- Peng F, Setyawati MI, Tee JK, Ding X, Wang J, et al. Nanoparticles promote in vivo breast cancer cell intravasation and extravasation by inducing endothelial leakiness. *Nature Nanotechnology*, (2019); 14279-286.
- Aitte SA, Zain M. Study Of Changes In Blood Parameters And Calculation Of PCT , MPV And DPW For The Platelets Of Laboratory Females And Males Of Albino Mice During Exposure To Doses Of Pyrethroid Pesticide (Alphacypermethrin). (2019); 14(2): 71--78.
- Lodhi M, Shaikh S, Afridi A, Baig MT, Farooq L, et al. Comparison of Anti-Seizure Efficacy of Combined Extract of *Swertia chirata* and *Brassica nigra* with Standard Anti-Epileptic Drugs in PTZ Model. *Health ...*, (2019); 8160-167.
- Lange ME, Uwiera RRE, Inglis GD. Housing Gnotobiotic Mice in Conventional Animal Facilities. *Current Protocols in Mouse Biology*, (2019); 9e59.
- Suárez AI, Chávez K (2018) Appraisal of Medicinal Plants with Anticancer Properties in South America. *Anticancer plants: Properties and Application*. Singapore: Springer Singapore. pp. 229-283.
- Rani R, Kansal VK. Study on cow ghee versus soybean oil on 7,12-dimethylbenz(a)-anthracene induced mammary carcinogenesis & expression of cyclooxygenase-2 & peroxisome proliferators activated receptor- γ in rats. *Indian Journal of Medical Research*, (2011); 133497-503.
- Kumar M, Dandapat S, Prasad Sinha M, Kumar A, Raipat BS. Different blood collection methods from rats: A review. *Balne Research Journal*, (2017); 846-50.
- Jagtap UB, Panaskar SN, Bapat VA. Evaluation of antioxidant capacity and phenol content in jackfruit (*Artocarpus heterophyllus* Lam.) fruit pulp. *Plant Foods Hum Nutr*, (2010); 65(2): 99-104.
- Almora Pinedo Y, Arroyo-Acevedo JL, Herrera-Calderon O, Chumpitaz Cerrate VM, Hañari Quispe R, et al. Preventive effect of *Oenothera rosea* on N-methyl-N-nitrosourea- (NMU) induced gastric cancer in rats. *Clinical and Experimental Gastroenterology*, (2017); Volume 10327-332.
- Arroyo-Acevedo J, Herrera-Calderón O, Chávez-Asmat R, Anampa-Guzmán A, Chumpitaz-Cerrate V, et al. Protective effect of *Chuquiraga spinosa* extract on N -methyl-nitrosourea (NMU) induced prostate cancer in rats. *Prostate International*, (2017); 5(2): 47-52.
- Joshi S, Kumar S, Ponnusamy MP, Batra SK. Hypoxia-induced oxidative stress promotes MUC4 degradation via autophagy to enhance pancreatic cancer cells survival. *Oncogene*, (2016); 35(45): 5882-5892.
- Mier-Giraldo H, Díaz-Barrera LE, Delgado-Murcia LG, Valero-Valdivieso MF, Cález-Ramírez G. Cytotoxic and Immunomodulatory Potential Activity of *Physalis peruviana* Fruit Extracts on Cervical Cancer (HeLa) and Fibroblast (L929) Cells. *Journal of Evidence-Based Complementary & Alternative Medicine*, (2017); 22(4): 777-787.
- Rana AP, Kaur M, Zonunsanga B, Puri A, Kuka AS. Preoperative Peripheral Blood Count in Breast Carcinoma: Predictor of Prognosis or a Routine Test. *Int J Breast Cancer*, (2015); 2015964392.
- Akuru U. Effect of selected plants on Haematological parameters of DMBA-induced breast cancer of albino rats. *International Journal of Biochemistry & Physiology*, (2019); 4.
- Akinbami A, Popoola A, Adediran A, Dosunmu A, Oshinaike O, et al. Full blood count pattern of pre-chemotherapy breast cancer patients in Lagos, Nigeria. *Caspian J Intern Med*, (2013); 4(1): 574-579.
- Wm A, Dw N. Hematological Markers of In Vivo Toxicity. *Journal of Hematology & Thromboembolic Diseases*, (2016); 04(02).
- Oday Ali L. Study effect of Breast Cancer on Some Hematological and Biochemical Parameters in Babylon Province, Iraq. *IOSR Journal of Pharmacy and Biological Sciences (IOSR-JPBS)*, 9(3): 20-24.
- Zingue S, Njuh AN, Tueche AB, Tamsa J, Tchoupang EN, et al. In Vitro Cytotoxicity and In Vivo Antimammary Tumor Effects of the Hydroethanolic Extract of *Acacia seyal* (Mimosaceae) Stem Bark. *BioMed Research International*, (2018); 20182024602.
- Chen C-H, Wu S-H, Tseng Y-M, Liao J-B, Fu H-T, et al. Suppressive Effects of *Puerariae Radix* on the Breast Tumor Incidence in Rats Treated with DMBA. *Journal of Agricultural Science*, (2017); 9(7): 68-68.
- Rochet NM, Markovic SN, Porrata LF. The Role of Complete Blood Cell Count in Prognosis—Watch this Space! *Oral History Review*, (2012); 0876.
- Asuquo Etim E, Abiodun Emokpae M, Collins Ohwonigho A, Adebayo Yusuf A. Association of platelet count and platelet indices with stages of women breast cancer in Yola, Nigeria. *Hematology & Transfusion International Journal*, (2018); 6(1).



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License. To read the copy of this license please visit: <https://creativecommons.org/licenses/by-nc/4.0/>