



Full Length Research Article

Advancements in Life Sciences – International Quarterly Journal of Biological Sciences

ARTICLE INFO

Open Access



Date Received:
11/07/2023;
Date Revised:
02/12/2023;
Date Published Online:
31/12/2023;
Date Updated:
09/09/2025

Identification of therapeutic phytochemicals targeting B-cell lymphoma 2 (BCL2) as anti-acute myeloid leukemia agents: An in-silico approach

Author's Affiliation:

Department of Biochemistry,
College of Science, University of
Jeddah, Jeddah - Saudi Arabia

Nawal Helmi

Corresponding Author:

Nawal Helmi
Email:
nmhelmi@uj.edu.sa

How to Cite:

Helmi N (2023).
Identification of therapeutic
phytochemicals targeting B-
cell lymphoma 2 (BCL2) as
anti-acute myeloid leukemia
agents: An in-silico
approach. Adv. Life Sci.
10(4): 670-674.

Keywords:

Acute myeloid leukemia;
Cancer, BCL2;
Phytochemicals; Drug-like

Editorial Note:

You are viewing the latest
version of this article having
language corrections.

Abstract

Background: Acute myeloid leukemia (AML) is a deadly cancer. B cell lymphoma 2 (BCL2) is frequently upregulated in AML and plays a vital role in the viability of both AML and AML stem cells. This study aimed to identify novel phytochemicals against BCL2 and evaluate their pharmacokinetics and toxicity prediction using in-silico tools.

Methods: In-silico screening of phytochemicals against the BCL2 active site using the PyRx0.8 AutoDock tool, followed by in silico pharmacokinetic and toxicity predictions, was performed. Protein-protein interaction analysis was performed using the STRING database to assess the interactions between BCL2 and neighboring interacting proteins.

Results: In total, 1106 terpenoid compounds were screened to evaluate their binding affinity toward BCL2. Five natural compounds demonstrated strong binding to the BCL2 protein after extensive screening, detailed interaction analysis, and visual inspections. Notably, the top compounds exhibited binding energies comparable to or greater than the positive control (venetoclax). In addition, these compounds were found to bind to key BCL2 residues and possess good drug-like properties.

Conclusions: The identified phytochemicals represent an essential initial step in drug discovery for AML management. Experimental validation is required to optimize the identified phytochemicals as potential BCL2 inhibitors.



Introduction

Leukemia is a type of cancer that originates from hematopoietic stem cells or progenitor cells. It is a heterogeneous group of diseases that affects the normal development and function of blood cells. Acute myeloid leukemia (AML) is a particularly aggressive form of leukemia characterized by the abnormal growth and proliferation of myeloid progenitor cells. AML is a significant health problem worldwide, with a reported global incidence of over one million cases per year and a resulting mortality rate of more than 100,000 [1,2]. Chemotherapy is still the most commonly used primary treatment option for AML [3,4]. Despite this treatment option, patients with AML have a five-year survival rate of only approximately 28%, indicating poor outcomes. Globally, the overall survival rate for individuals with AML remains poor [5].

The B cell lymphoma 2 (BCL2) protein family, which is involved in the regulation of apoptotic cell death, has been linked to the pathogenesis and progression of various cancers [6,7]. In AML, BCL2 is commonly upregulated and plays a crucial role in promoting the survival of both AML and AML stem cells [8,9]. As BCL2 is a central regulator of the intrinsic apoptotic pathway involved in the development and progression of several malignancies, inhibition of this protein represents a promising therapeutic strategy for developing novel anticancer agents [10]. Venetoclax, also known as ABT-199, is a newly discovered small-molecule inhibitor that can be administered orally and has been designed specifically to target BCL2 [11]. While the majority of patients respond to venetoclax treatment, the extent and duration of this response remain suboptimal [12].

Terpenes, a class of organic compounds derived primarily from plants and trees, exhibit notable biological properties such as analgesic and anticonvulsant properties. Numerous studies have shown that certain terpenes can reduce inflammation symptoms by inhibiting pro-inflammatory cytokine release [13]. Terpenes also have antimicrobial, antiviral, antioxidant, and analgesic properties, aid in digestion, and have a variety of other biological activities. Their application ranges from influencing food health factors to acting as components in anti-cancer therapeutic modalities [14].

Biologists and chemists have widely adopted computer-aided drug design (CADD) as an essential component of a comprehensive drug discovery approach [15]. CADD is extensively utilized in the pharmaceutical industry to reduce costs and expedite the early-stage development of biologically novel and active compounds. Moreover, CADD is integral to drug discovery, design, and analysis processes [16]. Phytochemicals are plant-derived compounds that are often more chemically diverse and safer than

commercially available synthetic medications. Additionally, they usually possess a variety of pharmacological properties, including antibacterial, anticancer, antioxidative, and anti-inflammatory activities [17,18]. This study aimed to identify novel phytochemical hits, specifically terpenoid compounds, that can potentially be used as BCL2 inhibitors to combat AML.

Methods

Target protein retrieval and preparation

The crystal structure of the BCL2 protein in complex with venetoclax was retrieved from the PDB database (PDB ID: 6O0K) [19]. The first step in preparing the protein was to eliminate both water and co-crystallized ligand, which in this case was venetoclax. The structure was then saved in the .pdb format.

Protein-protein interaction (PPI) analysis

PPI analysis was performed using the STRING database for BCL2 and neighboring interacting proteins to gain a better understanding of the protein-interacting partners [20].

Phytochemical library preparation

In this study, the Plant Secondary Compound DataBase (PSC-db) [21] was used; a total of 1106 terpenoid compounds were obtained from this database. The compounds were downloaded in the .sdf format, minimized, and prepared for further screening.

Virtual screening (VS)

In this study, the screening process involved the utilization of PyRx0.8 software, which utilizes AutoDock Vina [22] to identify the binding between prepared phytochemicals (terpenoid compounds) and BCL2 protein. PyRx0.8 is an open-source VS application used in CADD techniques to screen libraries of compounds against the target protein. The ligand-protein complex exhibiting the highest negative binding energy was selected for further investigation, and the binding interactions were subsequently analyzed. The grid box of the BCL2 active site was set as X= -11.003, Y= 1.56, and Z= -9.00 for the screening of the prepared compounds.

Estimation of the physicochemical and drug-likeness properties

The physicochemical and drug-likeness properties of the 10 selected compounds were estimated using the SwissADME web server [23].

Results

This study included the analyses of PPI interaction of BCL2 and molecular docking-based VS of 1106 terpenoid compounds to identify potent natural

inhibitors of BCL2. PPI analysis revealed that BECN1, BAX, TP53, BAD, BCL2L11, BIK, BBC3, BID, BCL2L1, and FKBP8 were the closest proteins that interacted with BCL2.




Figure 1: Protein-protein interaction analysis. 10 neighboring proteins interacting with BCL2 (A), and k-means clustering of the network of interacting proteins (B).

Clustering analysis showed that these interacting proteins belonged to the following three distinct groups or clusters: BAX, BCL2, and BECN1 in one cluster; BAD, BBC3, BCL2L1, BCL2L11, BID, BIK, and TP53 in the second cluster; and FKBP8 alone in the third cluster (Figure 1 and Table 1).

cluster number	cluster color	gene count	protein name
1	Red	3	BAX
		3	BCL2
		3	BECN1
2	Green	7	BAD
		7	BBC3
		7	BCL2L1
		7	BCL2L11
		7	BID
		7	BIK
		7	TP53
3	Blue	1	FKBP8

Table 1: Clusters of 10 interacting proteins with BCL2.

Screening of the prepared library of 1106 terpenoid compounds, in which the compounds were mainly carotenoids and apocarotenoids, diterpenoids, hemiterpenoids, monoterpenoids, sesquiterpenoids, steroids, and triterpenoids, was performed by targeting the binding pocket of BCL2. In this study, venetoclax was used as a reference control for analyzing the binding patterns and positions of the docked compounds. Visual inspection was performed to assess the binding efficiencies of both venetoclax and the top-screened ligands. Based on the findings, the top ten terpenoid compounds were identified, as shown in Table 2 (Figure 2).

The physicochemical and drug-like properties of the selected compounds were evaluated, and nearly all compounds demonstrated drug-like characteristics, indicating their potential to be developed as drug molecules (Table 3).




Figure 2: Binding sites residues of the BCL2 protein and docked complex of selected terpenoids in its active site.

S. No.	Compound name	Binding energy (kcal/mol)
1.	Ginkgolide A	-10.12
2.	Gibberellin A34-catabolite	-10.09
3.	Gibberellin 1	-10.05
4.	Zexbrevin B	-9.95
5.	7beta,12alpha-Dihydroxykaurenolide	-9.92
6.	Caryoptin	-9.90
7.	Echinocystic acid	-9.88
8.	Enhydrin	-9.85
9.	Paucin	-9.81
10.	Isodonal	-9.79
11.	Venetoclax (control)	-9.79

Table 2: Binding energy of selected 10 terpenoids.

Based on the visual inspection of the binding positions, five compounds (Gibberellin 1, Zexbrevin B, Ginkgolide A, Phorbol, and 7beta,12alpha-Dihydroxykaurenolide) were selected for an in-depth interaction analysis. Gibberellin 1 was found to interact with the Leu137, Ala149, Arg146, Tyr108, Trp144, Gly145, Tyr202, Arg106, Asp103, Gln99, Ala100, Val148, Phe198, and Phe104 residues of BCL2. The Arg107 and Asp103 residues were involved in H-bonding, whereas the Leu137, Arg146, Trp144, Gly145, Tyr202, Arg106, Gln99, Ala100, Val148, and Phe198 residues interacted with Gibberellin 1 via van der Waals interactions (Figure 3A). Zexbrevin B interacted to the Gly145, Trp144, Tyr202, Phe198, Val148, Arg107, Asp103, Tyr108, Ala100, and Phe104 residues of BCL2. The Trp144, Phe198, Val148, Arg107, and Tyr108 residues participated in van der Waals interactions, whereas the Gly145 and Phe104 residues H-bonded with Zexbrevin B (Figure 3B). Ginkgolide A was found to interact with the Gly145, Tyr108, Val148, Tyr202, Ala100, Phe198, Asp103, Arg107, and Phe104 residues of BCL2. The Phe198, Arg107, and Phe104 residues participated in van der Waals interactions, whereas the Gly145, Tyr108, and Asp103 residues H-bonded with Ginkgolide A (Figure 3C). Phorbol interacted with the Asp103, Val148, Ala100, Gly101, Phe104, Gly145, Asn143, Trp144, Tyr108, Arg107, Phe198, and Tyr202 residues of BCL2. The Asp103 and Ala100 residues H-bonded with Phorbol, whereas the Val148, Gly101, Phe104, Gly145, Asn143, Trp144, Tyr108, Arg107, and Phe198 residues were involved in van der Waals interactions (Figure 3D).

Compound names	MW	formula	RB	HBA	HBD	lipinski	ghose	veber	egan	muegge	bioavail
Ginkgolide A	408.4	C20H24O9	1	9	2	0	0	0	0	0	0.55
Phorbol	364.43	C20H28O6	1	6	5	0	0	0	0	0	0.55
Zexbrevin B	364.39	C19H24O7	3	7	2	0	0	0	0	0	0.55
Gibberellin 1	348.39	C19H24O6	1	6	3	0	0	0	0	0	0.56
7beta,12alpha-Dihydroxykaurenolide	332.43	C20H28O4	0	4	2	0	0	0	0	0	0.55
Caryoptin	492.56	C26H36O9	8	9	0	0	2	0	0	0	0.55
Echinocystic acid	472.7	C30H48O4	1	4	3	1	3	0	1	1	0.56
Enhydrin	464.46	C23H28O10	7	10	0	0	0	0	0	0	0.55
Paucin	468.49	C25H32O10	5	10	3	0	0	1	1	0	0.55
Isodonol	404.45	C22H28O7	4	7	1	0	0	0	0	0	0.55

Table 3: Predicted physicochemical and Drug-likeness properties of selected 10 terpenoids.

In addition, 7beta,12alpha-Dihydroxykaurenolide interacted with the Tyr202, Trp144, Gly145, Ala100, Val148, Phe198, Asp103, Arg107, Phe104, and Tyr108 residues of BCL2. The Trp144, Arg107, Phe104, and Tyr108 residues participated in van der Waals interactions, whereas the Tyr202, Gly145, and Asp103 residues H-bonded with 7beta,12alpha-Dihydroxykaurenolide (Figure 3E).

Furthermore, the control compound venetoclax interacted with the Glu152, Phe153, Val133, Glu136, Val148, Asp103, Arg107, Ala100, Asn143, Trp188, Trp144, Asn192, Leu201, Tyr202, Gly145, Arg146, Tyr108, Phe104, Ala149, Phe112, Val156, Met115, Asp111, and Leu137 residues of BCL2 (Figure 3F).




Figure 3: Interacting residues of BCL2 protein with Gibberellin 1 (A), Zexbrevin B (B), Ginkgolide A (C), Phorbol (D), 7beta,12alpha-Dihydroxykaurenolide (E), and venetoclax (F).

Discussion

The BCL2 family of proteins plays a crucial role in intrinsic apoptosis. In AML, excessive expression of BCL2 proteins can contribute to apoptosis resistance and therapeutic failure. Therefore, investigating anti-apoptotic BCL2 inhibitors holds great promise for identifying novel pharmacological agents for cancer treatment [24]. This study screened a total of 1106 terpenoid compounds against the BCL2 protein. Among these compounds, Gibberellin 1, Zexbrevin B, Ginkgolide A, Phorbol, and 7beta,12alpha-Dihydroxykaurenolide demonstrated robust binding with BCL2. The Phe104, Met115, Leu137, Arg146, Val148, Ala149, and Phe153 residues of BCL2 have been reported to be important in inhibitor binding [25]. Interestingly, in this study, the hit compounds (Gibberellin 1, Zexbrevin B, Ginkgolide A, Phorbol, and

7beta,12alpha-Dihydroxykaurenolide) were observed to bind to these residues of BCL2. Venetoclax binds to Glu152, Phe153, Val133, Glu136, Val148, Asp103, Arg107, Ala100, Asn143, Trp188, Trp144, Asn192, Leu201, Tyr202, Gly145, Arg146, Tyr108, Phe104, Ala149, Phe112, Val156, Met115, Asp111, and Leu137 of BCL2. Interestingly, Gly145, Tyr108, Val148, Tyr202, Ala100, Asp103, and Phe104 were observed to be the common binding residues for interactions between BCL2 and the hits (Gibberellin 1, Zexbrevin B, Ginkgolide A, Phorbol, and 7beta,12alpha-Dihydroxykaurenolide) as well as venetoclax (Figure 3A-3F).

Binding energy serves as a determinant of the degree of interaction between a protein and ligand, and a higher (negative) value of this energy signifies the successful binding of the inhibitor to its target protein [26]. Interestingly, Gibberellin 1, Zexbrevin B, Ginkgolide A, Phorbol, and 7beta,12alpha-Dihydroxykaurenolide exhibited strong binding with BCL2, with higher binding energies than that observed for venetoclax, suggesting that these hits can be used as BCL2 inhibitors to combat AML.

Phytochemicals and their derivatives found in plants have shown promise for improving cancer treatment efficacy while reducing the adverse effects. Many of these biologically active compounds occur naturally and possess significant antitumor potential [27]. Developing an effective and adverse-effect-free anticancer therapy based on phytochemicals requires initial testing of natural plant extracts for their potential anticancer biological activity. This is followed by the purification of the active phytochemicals using bioassay-guided fractionation and further evaluation of their *in vitro* and *in vivo* effects. From 1940 to 2014, approximately half of the approved anticancer drugs originated from natural products or their derivatives [28].

This study screened phytochemicals, specifically terpenoid compounds, against the BCL2 protein. Gibberellin 1, Phorbol, Zexbrevin B, Ginkgolide A, and 7beta, 12alpha-Dihydroxykaurenolide were observed to bind strongly with BCL2, bind to key BCL2 residues, and have good drug-like properties. These phytochemicals have the potential to be used in the management of AML. However, further experimental

validations are required to optimize their usage as BCL2 inhibitors for AML management.

Acknowledgment

I want to thank my colleagues at the University of Jeddah for their feedback and support throughout the research process. In particular, I would like to thank Dr. Hanadi Baisa for her valuable insights and suggestions.

Conflicts of interest

The authors declare no conflict of interest.

References

1. Mortality GBD, Causes of Death C. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*, (2016); 388(10053): 1459-1544.
2. Disease GBD, Injury I, Prevalence C. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*, (2016); 388(10053): 1545-1602.
3. Dohner H, Estey E, Grimwade D, Amadori S, Appelbaum FR, et al. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood*, (2017); 129(4): 424-447.
4. Dohner H, Estey EH, Amadori S, Appelbaum FR, Buchner T, et al. Diagnosis and management of acute myeloid leukemia in adults: recommendations from an international expert panel, on behalf of the European LeukemiaNet. *Blood*, (2010); 115(3): 453-474.
5. Rowe JM. Perspectives on current survival and new developments in AML. *Best Practice & Research Clinical Haematology*, (2021); 34(1): 101248.
6. Belka C, Budach W. Anti-apoptotic Bcl-2 proteins: structure, function and relevance for radiation biology. *International Journal of Radiation Biology*, (2002); 78(8): 643-658.
7. Alam M, Ali S, Mohammad T, Hasan GM, Yadav DK, et al. B Cell Lymphoma 2: A Potential Therapeutic Target for Cancer Therapy. *International Journal of Molecular Sciences*, (2021); 22(19): 10442.
8. Carter BZ, Mak PY, Tao W, Warmoes M, Lorenzi PL, et al. Targeting MCL-1 dysregulates cell metabolism and leukemia-stroma interactions and resensitizes acute myeloid leukemia to BCL-2 inhibition. *Haematologica*, (2022); 107(1): 58-76.
9. Young AI, Timpson P, Gallego-Ortega D, Ormandy CJ, Oakes SR. Myeloid cell leukemia 1 (MCL-1), an unexpected modulator of protein kinase signaling during invasion. *Cell Adhesion & Migration*, (2018); 12(6): 513-523.
10. Kapoor I, Bodo J, Hill BT, Hsi ED, Almasan A. Targeting BCL-2 in B-cell malignancies and overcoming therapeutic resistance. *Cell Death & Disease*, (2020); 11(11): 941.
11. Scheffold A, Jebaraj BMC, Stilgenbauer S. Venetoclax: Targeting BCL2 in Hematological Cancers. *Recent Results in Cancer Research*, (2018); 212: 215-242.
12. Bohl SR, Bullinger L, Rucker FG. New Targeted Agents in Acute Myeloid Leukemia: New Hope on the Rise. *International Journal of Molecular Sciences*, (2019); 20(8): 1983.
13. Del Prado-Audelo ML, Cortes H, Caballero-Floran IH, Gonzalez-Torres M, Escutia-Guadarrama L, et al. Therapeutic applications of terpenes on inflammatory diseases. *Frontiers in Pharmacology*, (2021); 12: 704197.
14. Fan M, Yuan S, Li L, Zheng J, Zhao D, et al. Application of terpenoid compounds in food and pharmaceutical products. *Fermentation*, (2023); 9(2): 119.
15. Sabe VT, Ntombela T, Jhamba LA, Maguire GEM, Govender T, et al. Current trends in computer aided drug design and a highlight of drugs discovered via computational techniques: A review. *European Journal of Medicinal Chemistry*, (2021); 224: 113705.
16. Usha T, Shanmugarajan D, Goyal AK, Kumar CS, Middha SK. Recent Updates on Computer-aided Drug Discovery: Time for a Paradigm Shift. *Current Topics in Medicinal Chemistry*, (2017); 17(30): 3296-3307.
17. Newman DJ, Cragg GM. Natural Products as Sources of New Drugs over the Nearly Four Decades from 01/1981 to 09/2019. *Journal of Natural Products*, (2020); 83(5): 770-805.
18. Wali AF, Majid S, Rasool S, Shehada SB, Abdulkareem SK, et al. Natural products against cancer: Review on phytochemicals from marine sources in preventing cancer. *Saudi Pharmaceutical Journal*, (2019); 27(6): 767-777.
19. Birkinshaw RW, Gong JN, Luo CS, Lio D, White CA, et al. Structures of BCL-2 in complex with venetoclax reveal the molecular basis of resistance mutations. *Nature Communications*, (2019); 10(1): 2385.
20. von Mering C, Huynen M, Jaeggi D, Schmidt S, Bork P, et al. STRING: a database of predicted functional associations between proteins. *Nucleic Acids Research*, (2003); 31(1): 258-261.
21. Valdes-Jimenez A, Pena-Varas C, Borrego-Munoz P, Arrue L, Alegria-Arcos M, et al. PSC-db: A Structured and Searchable 3D-Database for Plant Secondary Compounds. *Molecules*, (2021); 26(4): 1124.
22. Dallakyan S, Olson AJ. Small-molecule library screening by docking with PyRx. *Methods in Molecular Biology*, (2015); 1263: 243-250.
23. Daina A, Michielin O, Zoete V. SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Scientific Reports*, (2017); 7: 42717.
24. Wei Y, Cao Y, Sun R, Cheng L, Xiong X, et al. Targeting Bcl-2 Proteins in Acute Myeloid Leukemia. *Frontiers in Oncology*, (2020); 10: 584974.
25. Bultum LE, Tolossa GB, Lee D. Combining empirical knowledge, in silico molecular docking and ADMET profiling to identify therapeutic phytochemicals from Brucea antidysentrica for acute myeloid leukemia. *PLoS One*, (2022); 17(7): e0270050.
26. Meng XY, Zhang HX, Mezei M, Cui M. Molecular docking: a powerful approach for structure-based drug discovery. *Current Computer-Aided Drug Design*, (2011); 7(2): 146-157.
27. Choudhari AS, Mandave PC, Deshpande M, Ranjekar P, Prakash O. Phytochemicals in Cancer Treatment: From Preclinical Studies to Clinical Practice. *Frontiers in Pharmacology*, (2019); 10: 1614.
28. Newman DJ, Cragg GM. Natural Products as Sources of New Drugs from 1981 to 2014. *Journal of Natural Products*, (2016); 79(3): 629-661.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License. To read the copy of this license please visit: <https://creativecommons.org/licenses/by-nc/4.0/>