

ARTICLE INFO

Open Access



Date Received:
23/08/2023;
Date Revised:
09/08/2024;
Available Online:
31/12/2024;

The protective role of ethanolic extract of ginseng (Panax ginseng) against adverse physiological and histological alterations in the liver of female rabbits exposed to chlorine

Shireen Ali Hasan*, Alyaa Abdalrazaq Abass, Layth Jabbar

Author's Affiliation:

Department of Pharmacology and Toxicology, College of Pharmacy, University of Thi-Qar, Thi-Qar – 64001 – Iraq

***Corresponding Author:**

Shireen Ali Hasan
Email:
Shireenalihasan@utq.edu.iq

How to Cite:

Hasan SA, Abass AA, Jabbar L (2025). The protective role of ethanolic extract of ginseng (Panax ginseng) against adverse physiological and histological alterations in the liver of female rabbits exposed to chlorine. *Adv. Life Sci.* 12(1): 185-190.

Keywords:

Ginseng; Chlorine; MDA; Liver enzymes; Blood parameters

Editorial Note:

You are viewing the latest version of this article having minor corrections related to the use of English language and in references section.

Abstract

Background: Ginseng (*Panax ginseng*) has garnered considerable attention due to its medicinal properties. This study aimed to assess the protective effects of ginseng ethanolic extract against chlorine-induced toxicity in female rabbits.

Methods: Thirty-two female rabbits were randomly divided into four groups: a negative control group receiving chlorine-free water, a positive control group receiving tap water with chlorine (0.05 ppm/L), a group exposed to chlorine (5 ppm/L) in water, and a group exposed to chlorine (5 ppm/L) along with ethanolic ginseng extract (200 mg/kg body weight). Treatment lasted for four weeks.

Results: revealed that oral exposure to chlorine led to oxidative stress, evidenced by elevated levels of malondialdehyde (MDA) and liver enzymes (AST & ALT), as well as suppressed blood parameters (PCV, HB, and RBC count), and histological liver alterations compared to the normal control group. Conversely, administration of ginseng extracts alongside chlorine for four weeks reduced MDA levels and liver enzyme activity while enhancing blood parameters, accompanied by decreased inflammatory histological changes.

Conclusion: These results indicate that ginseng extract has strong antioxidant activity, reducing the toxicity and related pathophysiological effects of chlorine.

Introduction

Chlorine (Cl₂) is extensively used as a water disinfectant because of its ability to kill waterborne diseases. Nonetheless, its use is associated with the generation of disinfection by-products (DBPs) and possible ill effects on health [1]. Chlorine is pivotal in developed nations for water sanitation for effectively combating waterborne diseases, including cholera, typhoid, and dysentery [2]. Nonetheless, its volatile nature creates inhalation hazards, and its reaction with organic contaminants in water produces toxic disinfection by-products (DBPs). The DBPs comprising the trihalomethane, chloroform, bromoform, haloacetic acid, and others are known to be toxic to health and physiology and even lead to the development of cancer. The trihalomethanes and haloacetic acids are usual contaminants in treated water and are tracked in various countries for the measurement of DBP exposure and the resultant negative impact on the functioning of the body [3]. Additionally, exposure to chlorine and its derivatives triggers oxidative stress, causes disruption of the metabolic pathways, and causes malfunction of various organs. In effect, chlorine as a water disinfectant is vital but poses possible ill effects on health through DBP generation and the initiation of oxidative stress, with the use of appropriate conditions and alternative methods for safe water requiring careful management [4,5]. Ginseng is a commonly known medicinal plant with various bioactive molecules and natural free radicals. The compound has significant strength in strengthening the endogenous antioxidant defense against reactive oxygen species (ROS), restoring the optimal balance, and preventing disease [6]. Of the thousands of medicinal herbs, *Panax ginseng* is among the most economically important. *Panax ginseng* has been used for centuries in America for medicinal and nutritional purposes, with the roots and leaves being edible [6]. *Panax ginseng*, commonly called the Korean ginseng or the Asian ginseng, is mainly located in northeast Asia, including countries like China, Korea, Bhutan, and eastern parts of Siberia [7]. Ginsenosides (ginseng saponins), the active molecules in ginseng, along with the phenolic molecules, polysaccharides, and polyacetylenes, are the molecules responsible for the pharmacological activity [8]. The pharmacological activities include antioxidant activity, immune enhancement, anti-aging effects, reduction of stress, anti-inflammation approach, anti-neoplastic effect, and anti-tumor activity. Considering the multifunctional remedial utility of ginseng, the present study intended to evaluate the protective activity of ginseng ethanolic extract against the deleterious physiological outcomes caused by chlorine exposure in female rabbits [9].

The research endeavors to utilize the antioxidant activity of ginseng against the oxidative stress and other deleterious outcomes that arise following chlorine exposure, providing hope for therapeutic applications in reducing chlorine toxicity.

Methods

Procurement of *Panax Ginseng*:

The roots of *Panax ginseng* were purchased from the marketplace of Thi Qar city, Iraq.

Preparation of Roots

Washed the ginseng roots with distilled water to eliminate any contaminants. Next, dried the roots slowly at room temperature for 5-7 days in the shade until fully dry.

Cleaning and Grinding

Cleaned the dried roots to remove any remaining debris or dirt. Grind the cleaned roots into a fine powder using a grinder.

Extraction of Leaves:

50g of the powdered ginseng leaves were taken. This powder was measured and stored in a round-bottom flask.

Preparation of Extraction Solvent

The round-bottom flask with powdered ginseng leaves was treated with 200 mL of 70% ethanol.

Extraction Process

The mixture stood for 12 hours in a reflux extractor setup. This facilitated the effective extraction of bioactive compounds from ginseng leaves into the ethanol solvent.

Filtration

The mixture was filtered after the extraction period with Whatman filter paper (No. 31) in order to remove the extracted solution from the solid residue. The process facilitated the acquisition of a clear extract.

Drying of Extract

The filtered extract was then allowed to dry naturally in the shade in a petri dish that had been cleaned beforehand. By this method, the ethanol solvent in the extract evaporated, and what remained was the dried ginseng extract.

Collection and Storage

After drying, the ginseng extract is gathered and kept in a tightly sealed container at 4°C for future use. Adequate storage retains the stability and strength of the extract in the course of time.

Dosage Preparation

The desired dosage for the experiment was determined. In this study, doses of 100 mg/kg and 200 mg/kg of the ginseng extract were utilized.

Usage in Experiment

The prepared ginseng extract was then used in the experiment, as described in the methodology, to evaluate its protective efficacy against chlorine-induced adverse effects in female rabbits [10]. Treatment and Animal Housing Conditions: Thirty-two female rabbits, each weighing approximately (grams), were housed in the departmental Animal House in pairs within plastic cages. The animals were acclimated for 14 days prior to the experiment under controlled conditions (temperature: 25 ± 2 °C, 12-hour light/dark cycle) with free access to food and water. The rabbits were randomly divided into four experimental groups, with eight rabbits per group, as follows:

- Group 1 (Negative Control): Rabbits received chlorine-free drinking water.
- Group 2 (Positive Control): Rabbits received tap water with chlorine estimated at 0.05 ppm/L.
- Group 3: Rabbits received water with chlorine at a concentration of 5 ppm/L.
- Group 4: Rabbits received water with chlorine at 5 ppm/L, along with ethanolic ginseng extract (200 mg/kg body weight), administered one hour after water intake.

According to Gabry et al. (2014), treatment continued for four weeks [11].

Blood samples, traits, and physiological assay:

Hemoglobin (Hb) Quantification: Spectrophotometric determination of the hemoglobin concentration has been achieved through a method described in reference [12]. **Packed Cell Volume (PCV):** Packed cell volume, or hematocrit value, was determined through a spectrophotometric method following the procedure described in reference [12]. **Red Blood Cell (RBC) Count:** Red blood cell count was measured through spectrophotometry following the procedure described in reference [12]. **Malondialdehyde (MDA) Measurement:** The levels of malondialdehyde, the oxidative stress marker, were determined with the help of the Enzyme-Linked Immunosorbent Assay (ELISA) method as described in reference [13]. **Transaminase Enzymes (AST, ALT) and Alkaline Phosphatase:** Transaminase enzymes (aspartate transaminase - AST, alanine transaminase - ALT) and alkaline phosphatase levels were determined with a Reflotron apparatus and with commercially available kits following the manufacturer's guidelines, as discussed in [14]. The above analysis methods were selected in order to examine several physiological parameters, such as

blood characteristics and oxidative stress markers, in order to examine the effect of ginseng ethanolic extract against the toxic effects of chlorine exposure in female rabbits.

Tissue Samples: liver tissues were collected from rabbits, washed with normal saline, prepared then fixed by using 10% formalin for histological examination.

Chemical implementation:

Chlorinated water at concentration (5 ppm/L) prepared previously before each use for animal groups was treated with chlorinated water [15]. Tap water quality criteria are investigated to assess the liberated chlorine levels (ppm/L) by gathering samples of water and using a pocket digital colorimeter that analyzes water samples for liberated chlorine by following the instructions for use. Likewise, the device is modern technology with high sensitivity, having a range from 0.01 up to 1.00 ppm/L [16]. The Statistical Analysis: Mean values of serum indices were analyzed by one-way analysis of variation (ANOVA), obtained mean differences and standard deviations (Mean \pm SD) between treated and control groups, P-values ($P < 0.05$) are considered statistically significant. Table 1 serum MDA and blood parameters in control group and treated groups with Ginseng extract plus chlorine in female rabbits. Data are expressed as mean \pm SD (n=8). The different letters refer to significant difference ($P < 0.05$).

Results

The results of Table 1 indicate that the exposure to tap water 2nd group and Chlorine 5 ppm water 3rd group induced remarkable alterations in blood indices. There is a significant reduction, $p \leq 0.05$, in the average of RBC, Hb, and PCV compared with control 1st group. While the treatment with ethanolic extract of ginseng and chlorine resulted in a notable increase in RBC, Hb, and PCV values, comparison with group exposed to chlorine alone shows that the decline in these components was mostly observed in the second and third groups. Also, the data illustrated in the Table 1 revealed remarkable elevation in MDA level to Tap water (2nd group) and Chlorine 5 ppm water (3rd group) compared to normal rabbits, whereas the receiving animals ethanolic extract ginseng and chlorine (group 4th) prominent reduction in MDA level as compared to the chlorine group.

Data obtained from the Table 2 exhibited that the oral exposure of Tap water (2nd group) and Chlorine 5 ppm water (3rd group) induced prominent increase $p \leq 0.05$ in levels of liver enzymes (AST and ALT) compared to the corresponding control, whereas the animals received ethanolic extract ginseng and chlorine (group 4th) produced a notable decrease in

levels of the above enzyme's comparison to chlorine. These results clarified that the hepatotoxicity was induced in normal, healthy rabbits with metabolic activity stable, so any effect of chlorine or the indirect influence of liver enzymes on hepatocytes' activity.

Parameters Groups	RBC (10 ⁶ cell/mm ³)	Hb g/100ml	PCV%	MDA μ mol/L
Chlorine free water 1st group	6.76±0.35a	12.94±0.57 a	43.78±2.14 a	0.54±0.03 b
Tap water 2nd group	4.87±1.21b	10.87±0.36 b	36.16±2.12 b	2.18 ±0.10 a
Chlorine 5ppm water 3rd group	3.92±0.34 b	9.01±1.08 c	26.89±1.99 c	3.13 ±1.86 a

Table 1: Serum MDA and blood parameters in control group and treated groups with Ginseng extract plus chlorine in female rabbits. Data are expressed as mean ± SD (n=8). The different letters refer to significant difference (P< 0.05).

Parameters Groups	ALT unit/L	AST unit/L
Chlorine free water (1st group)	52.9 ±2.68c	38.17±3.29c
Tap water (2nd group)	74.87±1.21b	57.87±0.36 b
Chlorine 5ppm water 3rd group	83.92±0.34a	72.01±1.08 a
Chlorine 5ppm water GN(200mg/kg) 4th group	54.81±0.22 c	45.82±0.83c

Table 2: liver enzymes in control group and treated groups with Ginseng extract plus chlorine in female rabbits. Data are expressed as mean ± SD (n=8). The different letters refer to significant difference (P< 0.05).




Figure 1: (A, B, C, and D): Liver section of female rabbits (control group), the normal structure of the hepatic lobule with normal central vein (CV) and portal vein (PV). (Fig B and C) Female rabbits treated with chlorine for four weeks, severe congestion and dilatation in portal vein, and sinusoid extension (thin arrow), with fibrosis, aggregation of inflammatory cells around bile duct and vein (thick arrow), and hyperplasia of bile duct (BD). H&E. X40 I. (Fig: D) female rabbits treated with ginseng extract (200mg/kg B.W) and chlorine for 4 weeks, sinusoid less dilated (thick arrows) with regeneration in some area which appeared with early hyperplasia (head arrow), inflammatory cells filtration (thin arrow) and less congested in central vein. 40 X H&E.

Discussion

As shown in table (1) the results indicate that the exposure to Tap water 2nd group and Chlorine 5ppm water 3rd group induced remarkable alteration in blood indices, there is a significant reduction $p < 0.05$ in the average of RBC, Hb and PCV compared with control 1st group, While the treatment with ethanolic extract ginseng and chlorine resulted in notable increase in RBC, Hb and PCV values comparison with group exposed to chlorine alone the decline in these components was mostly observed in the second and third groups could attributed to the high portions of chlorine and due to disinfection the water by chlorine and overlapping or suppression the excretion of erythropoietin hormone from the kidney and with a low portions from the liver, which the first and most responsible for synthesis of erythrocytes from bone marrow, may be due to hematotoxicity caused by chlorine treatment which lead to decrease synthesis of red blood cells in bone marrow. Chlorine caused a significant decrease in the levels of RBC count, Hb and PCV, indicating anemia. This anemia may come in part from the effects of free radicals generated by the chlorine on RBC. Several studies have suggested that RBC are particularly vulnerable to oxidative stress because they are exposed to higher concentrations of oxygen and, thus, Hb can easily be oxidized [17]. On the other hand, the activity of ALAD (d-aminolevulinic acid dehydratase which represents a key regulatory enzyme of heme synthesis pathway) is highly sensitive to the presence of stressful chemical; ALAD is a sulfhydryl-containing and its inhibition can be attributed to the binding of chlorine with sulfhydryl groups), the inhibition of ALAD enzyme and the reduction of Hb and RBC led to decreased heme synthesis and ultimately anemia and therefore lowering in Hb, PCV estimates this agreed and confirmed with [1,18], increased in estimates of blood traits Hb and PCV table (1) in the treated with ethanolic extract ginseng and chlorine group could be attributed to The extract of *P. ginseng* possesses antioxidant properties was explained by the presence of ginsenosides which are the main active compounds in ginseng, that reducing oxidative stress [18]. This protective effect may be due to the higher ability to eliminate free radicals and protect the cell membranes from the lipid peroxidation [19], The data illustrated in table (1) revealed remarkable elevation in MDA level to Tap water(2nd group) and Chlorine 5ppmwater (3rd group) compared to normal rabbits, whereas the receiving animals ethanolic extract ginseng and chlorine (group 4th) prominent reduction in MDA level as compared to the chlorine group.

The outcomes of this study (Table 1) revealed significantly ($p < 0.05$) higher in malondialdehyde

concentration especially in female rabbit serum that treated Tap water and Chlorine 5ppmwater compared with another groups could be attributed to the portability of (chlorine) to activating the oxidation and producing the free radicals and increasing fat peroxidation leading to oxidative stress, therefore increasing malondialdehyde concentration as an important production of peroxidation or (chlorine) may be cause activating to enzymes that in change of unsaturated fatty acid oxidation, which lead to producing hydrogen peroxide and therefore disorder in balance between production of free radicals and antioxidant defenses which caused tissues damage and increased MDA production, this outcomes agreed with [20,21]. The present study showed that the co-administration of ginseng combined with chlorine may have beneficial role and protective effects against chlorine intoxication. That role played by ginsenosides free phenolic fractions have been shown to induce the cytosolic antioxidant enzyme superoxide dismutase by enhanced nuclear protein binding to its gene regulatory sequences. Ginseng contains polyphenols, flavonoids, saponins, these properties effects might be due to the potential antioxidant, free radical scavenging radical mechanism and detoxification effected, these results similar to those [7, 9]. Data obtained from table (2) the outcomes exhibited that the oral exposure of Tap water (2nd group) and Chlorine 5ppmwater (3rd group) induced prominent increase $p \leq 0.05$ in levels of liver enzymes (AST and ALT) compared to the corresponding control, whereas the animals received ethanolic extract ginseng and chlorine (group 4th) produced a notable decrease in levels of the above enzymes comparison to chlorine. The progressively increased in the outcomes of (AST and ALT) in second and third groups could be attributed to the oxidative stress in liver and the injury done to tissues that contains those enzymes especially liver cells resulting in production free radicals (ROS) that impact cellular activity especially cell wall functions, as considered chlorine as toxic and harmful impact materials to cell functions from through the elevation in the biomarkers of liver (Transaminases activity) indicator to cellular damage of liver hence released these enzymes in to circulation, the oxidative stress and hepatic injury induced by this by- products, similar results were agreed and confirmed by some researches [22]. On the other hand, co-administration of ginseng extract combined with chlorine restored the levels of enzymes in the serum of the rabbits as an indication of protective effect of ginseng extract against liver damage induced by chlorine. This is consistent with [26], who stated that ginseng has a productive effect on hepatotoxicity. The hepatoprotective effects of ginseng extract by diminishing AST and ALT activities

in serum due to three of its components (falcarinol, octadecadienoic acid and acetate [23,24].

Histopathological Study of Liver

A histopathological assessment in the liver of the control groups were evaluated, consist of a repeated hexagonal unit called hepatic lobules, each lobule is occupied by center vein, hepatocytes and sinusoid are arranged in cords radiating from the central hepatic (Fig. 1A) the inter lobular in the portal vein where the branches of a hepatic artery, hepatic portal vein and bile ducts, blood sinusoids were lined with Kupffer cells and endothelial cells (Fig. 1A).

After 4 weeks from orally given chlorine, the liver of rabbits showed alteration in structure, dilation and congestion of central vein and sinusoid capillaries, moderate enlargement of hepatic cell, which containing to the portal triad to the portal triad could be revealed to the formation of fibrosis, infiltration of mononuclear inflammatory cells in the portal area and central vein, thickening in the lining endothelial layer, some nuclei chromatin was fragmented and cytoplasm contained many vacuoles and necrotic in some area (Fig 1B,1C). When rabbits treated with ginseng extract (200mg/kg) plus chlorine the liver examination revealed amelioration section from treated chlorine 4 weeks alone clear central and portal veins were filled with erythrocytes, all hepatocytes had eosinophilic cytoplasm, some of these vacuolated cytoplasm and sinusoid were narrowed and irregular shaped (Fig 1D.), This results clarified as the hepatotoxicity was artificially in normal, healthy rabbits with metabolic activity stable, so any effect of chlorine or the indirect influence of liver enzymes on hepatocytes activity. Previous suggest on showed clear histological changes included damaged of the hepatocytes, wider sinusoid spaces and presence of binucleated cells indicating proliferation and regeneration. inflammatory cells in the hepatic tissue and necrosis due to chlorine exposure may suggest that chlorine might interact with proteins and enzymes of liver interstitial tissue interfering with the antioxidant defense mechanism and leading to generation ROS, which in turn may imitate an inflammatory. The ameliorating effects of ginseng which significantly inhibits liver fibrosis by inhibiting activation, proliferation and expression of collagen, transforming growth factor and tissue inhibitor of metalloproteinase-1 in hepatic stellate cell, the major cause of liver fibrosis [20-24]. However, the ginseng extract regulating inducible hepatic enzymes, and might be associated with modulating liver cytochrome P450 activation and protein 152 phosphorylation, similar have also been reported by [6].

Author Contributions

Shireen Ali Hasan conceptualized, supervised the study, and collected data, performed experiments, did data analysis and reviewing manuscript. Alyaa Abdalazraq Abass helped in computational analysis and Layth Jabbar was involved in proofreading the manuscript

This study used the procedures agreed upon by the Standing Committee for Scientific Research Ethics at University of Thi-Qar/ College of Pharmacy.

References

- Nickmilder M, Bernard A. Ecological association between childhood asthma and availability of indoor chlorinated swimming pools in Europe. *Occupational and environmental medicine*, (2007); 64(1): 37-46.
- Lantagne DS, Cardinali F, Blount BC. Disinfection by-product formation and mitigation strategies in point-of-use chlorination with sodium dichloroisocyanurate in Tanzania. *The American Journal of Tropical Medicine and Hygiene*, (2010); 83(1): 135.
- Font-Ribera L, Kogevinas M, Zock J-P, Nieuwenhuijsen MJ, Heederik D, et al. Swimming pool attendance and risk of asthma and allergic symptoms in children. *European Respiratory Journal*, (2009); 34(6): 1304-1310.
- Bawa G, Afolayan S, Olumeyan D, Ashiru R. Effects of various durations of water deprivation on performance of weaner rabbits in a Sub-Humid environment. *Pakistan Journal of Nutrition*, (2006); 5(6): 551-554.
- Burch JB, Everson TM, Seth RK, Wirth MD, Chatterjee S. Trihalomethane exposure and biomonitoring for the liver injury indicator, alanine aminotransferase, in the United States population (NHANES 1999–2006). *Science of The Total Environment*, (2015); 521: 226-234.
- Ramesh T, Kim S-W, Sung J-H, Hwang S-Y, Sohn S-H, et al. Effect of fermented *Panax ginseng* extract (GINST) on oxidative stress and antioxidant activities in major organs of aged rats. *Experimental Gerontology*, (2012); 47(1): 77-84.
- Morsy FA. Protective effect of ginseng against toxic effect of ochratoxin (OA) in kidney of male rats. *The Egyptian Journal of Hospital Medicine*, (2002); 7(1): 151-167.
- Kim H, Hong M-K, Choi H, Moon H-S, Lee H-J. Chemopreventive effects of korean red ginseng extract on rat hepatocarcinogenesis. *Journal of Cancer*, (2015); 6(1): 1.
- Diab AE-AA, Hendawy AA, Zahra MH, Hamza RZ, Abd El-Aziz E, et al. Ameliorative role and antioxidant effect of propolis and ginseng against reproductive toxicity of chlorpyrifos and profenofos in male rats. *Marsland press*, (2012); 9: 2557-2567.
- Abbasi Oshaghi E, Khodadadi I, Saidijam M, Yadegarazari R, Shabab N, et al. Lipid lowering effects of hydroalcoholic extract of anethum graveolens L. and dill tablet in high cholesterol fed hamsters. *Cholesterol*, (2015); 2015(1): 958560.
- Gabry MS, Kader DHA, Moustafa M, Elenany AAH. Effect of some antioxidants on the prostate of adult and aged albino rats: a histological and immunohistochemical study. *Journal of Applied Pharmaceutical Science*, (2014); 4(2): 017-026.
- Ahemen T, Abu A, Anongu T. The effect of feeding varying levels of water spinach (*Ipomoea aquatica*) leaf meal on reproductive tract morphometry, haematology and serum biochemical parameters of female rabbits. *Middle-East Journal of Scientific Research*, (2015); 23(11): 2791-2796.
- Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analytical biochemistry*, (1979); 95(2): 351-358.
- Burtis CA, Ashwood ER. *Tietz textbook of clinical chemistry*. Philadelphia, (1999); 1654-1655.
- Hocking MB *Handbook of chemical technology and pollution control*. Chapter: Book Name. 2006 of publication; Elsevier.
- Jeong CH, Postigo C, Richardson SD, Simmons JE, Kimura SY, et al. Occurrence and comparative toxicity of haloacetaldehyde disinfection byproducts in drinking water. *Environmental science & technology*, (2015); 49(23): 13749-13759.
- Özkan C, Kaya A, Akgül Y. Normal values of haematological and some biochemical parameters in serum and urine of New Zealand White rabbits. *World rabbit science*, (2012); 20(4): 253-259.
- Al-Zwean DH. The effect of water treated with potassium permanganate (KMnO₄) and chlorine on some hygienic and physiological parameters in the local rabbits. *Journal of Entomology and Zoology Studies*, (2017); 5(3): 1554-1559.
- Chung I-M, Kim J-W, Seguin P, Jun Y-M, Kim S-H. Ginsenosides and phenolics in fresh and processed Korean ginseng (*Panax ginseng* CA Meyer): Effects of cultivation location, year, and storage period. *Food chemistry*, (2012); 130(1): 73-83.
- Alfanie I, Muhyi R, Suhartono E. Effect of heavy metal on malondialdehyde and advanced oxidation protein products concentration a focus on arsenic, cadmium, and mercury. *J Med and Bioengineering*, (2015); 4.
- Bernard A, Voisin C, Sardella A. Con: respiratory risks associated with chlorinated swimming pools: a complex pattern of exposure and effects. *American journal of respiratory and critical care medicine*, (2011); 183(5): 570-572.
- Ali SA, Al-Derawi KH, Al-Mansour NAA. Toxic effects of lead-induced Damage in liver of rats male and the role of ethanolic ginseng extract as protective agents (histological and physiological). *Merit Research Journal of Medicine and Medical Sciences*, (2018); 9: 307-315.
- Huu Tung N, Uto T, Morinaga O, Kim YH, Shoyama Y. Pharmacological effects of ginseng on liver functions and diseases: a minireview. *Evidence-Based Complementary and Alternative Medicine*, (2012); 2012(1): 173297.
- Voces J, Cabral de Oliveira A, Prieto JG, Vila L, Perez AdC, et al. Ginseng administration protects skeletal muscle from oxidative stress induced by acute exercise in rats. *Brazilian journal of medical and biological research*, (2004); 37: 1863-1871.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License. To read the copy of this license please visit: <https://creativecommons.org/licenses/by-nc/4.0/>